

Dermatologic Parasitology

Scabies

(The itch disease)

Scabies is an infestation by the mite *Sarcoptes scabiei* which is a very small microorganism with dimensions of 0.4 x 0.3 mm for the female and 0.2 x 0.15 for the male. It appears by the magnifying lenses as a very small dot having many legs with multiple joints for each leg.

Life cycle:

When come in contact with the human skin, the fertilized female mite attack (and dig in) the stratum corneum making a tunnel (called burrow) in which it lay eggs. Each fertilized female lay 3 eggs per day, the eggs hatch in 3 days into male and female larvae, that need 3 weeks to become adult mites. The life span of the mite is 6 weeks but it die in 3 days if get out of human body. The new females are fertilized by males and then either remain in the burrow or leave it to form other burrow in the same patient or to infest another person to repeat the cycle.

Source: Prisons, hotels, institutes, or military places.

Transmission: close skin-to-skin contact as with sex partner, children playing, or health care worker. Mites can remain alive for over 2 days on clothing or in bedding, and therefore, scabies can be acquired by sharing beds or clothing.

Clinical features: If a patient receive the mite, symptoms will not appear before 3 weeks when immune response reaches its peak, but if he was previously infested, he will react after 3 days.

1- Generalized disturbing nocturnal itch that is aggravated by heating is the most important symptom.

2- Primary lesions are the burrows and vesicles. The burrow is a zigzag (S or C-shaped) skin colored elevated line or ridge that ends with a vesicle. Common sites of burrows are the finger webs, sides of fingers, axillae, umbilicus, buttocks, male genitalia, female breast and areola, and, in infants, the palms and soles

3- The secondary lesions appear as allergic papules and urticaria, scratch marks, and secondary bacterial infections.

4- Nodular lesions develop in 10 % of patients (especially in the genitalia and axillae) that may persist for long duration (3-4 months) after successful eradication of the mite, and if so, they are called post-scabetic nodules.

Diagnosis: Clinical findings, confirmed, if possible, by microscopy.

- Clinically

Itching Generalized
 Nocturnal
 ↑ by heat

Family member or close contact affection

Site of lesions

- Finding the borrow and demonstration of the mite, its eggs or even just the feces will confirm the diagnosis.
- Sometimes when the mite cannot be demonstrated, a “therapeutic test” will clinch the diagnosis.

Treatment

- Scabicides: should be applied to the entire body from neck to feet, being sure to treat the genitals, umbilicus, and all body folds.
- 1- Gamma benzene hexachloride (lindane): is applied only once and is very effective. It may be toxic to infants.
 - 2- Permethrin : effective, non-toxic. it is the treatment of choice in young children and pregnant women
 - 3- Benzyl benzoate is applied daily for 3 days.
 - 4- Sulfur 5-10% in petrolatum: applied for 3 days, simple, effective and cheap but messy and irritant.
 - 5- Crotamiton: twice a day for 3-5 days, 25%. failure rate but safe in infants and during pregnancy.
 - Treatment of itching: the worst itching fades in 1-2 days but sedating antihistamines and corticosteroids may be needed.
 - Treatment of secondary bacterial infection.
 - Treatment of contacts.
 - Environmental treatment

Pediculosis

It is infestation of human with lice. A louse is a small (2-4mm in length) wingless flattened insect (6 legged).

There are 3 common types:

- 1- The head louse: *Pediculus humanus capitis*
- 2- The body louse: *Pediculus humanus corporis*
- 3- the pubic louse: *Phthirus pubis*

Life cycle:

The cycle starts when the adult female lays 8 eggs/day on the surface of skin (where temperature is optimum for the eggs). These hatch in 8 days into nymphs which require another 8 days to become mature males and females. The mature adult female lives about one month.

Pediculosis Capitis

Infestation of the scalp by head louse, It is more common in school and pre school age but no age is immune.

Transmitted by:

- Direct head-to-head contact in schools classes, small rooms, especially in winter.
- Indirect contact by shared hats, caps, brushes, combs.

Clinical Features:

- 1- Pruritus and itching that is persistent day and night especially in the occipital region and sides of neck
- 2- Eczematisation of the neck.
- 3- Secondary bacterial infections due to scratching resulting in exudation, crusting and impetigo.
- 4- cervical lymphadenopathy.

Diagnosis

Clinical findings, confirmed by detection of nits and/or lice. Finding a louse is difficult. Majority of patients have a population of < 12 lice. Mainly we find the nits (eggs), these vary in number from only a few to thousands *Nits* should be differentiated from the *scales* (dandruff) that attach the scalp hair.

Nits	Scales
<i>Nits</i> are firmly attached to hair shaft and difficult to be removed.	<i>Scales</i> are loosely attached and can easily be removed.
<i>Nits</i> are regular in shape and form an angle of 45 with the hair shaft.	<i>Scales</i> are irregular in shape and encircle the hair shaft or lie free between them.
When you press the <i>nit</i> between two nails, you will hear a click sound (i.e. click sign positive)	Click sign is negative with the <i>scale</i> .
<i>Nits</i> illuminate and appear shiny and white by wood's light examination because they are viable.	<i>Scales</i> are not viable and will not appear shiny.
By light microscopic examination we can see the embryo inside the <i>Nit</i> .	Scale does not contain embryo.

Treatment

- *Pediculocides*

- 1- Permethrin shampoo is worked into the scalp, left on for 5-10 minutes and rinsed out.
- 2- lindane (Gamma benzene hexachloride) shampoo.
- 3- Malathion
- 4- benzyl benzoate
- 5- pyrethrin shampoo
- 6- Kerosine: applied for 3 successive days, it may cause irritation.
 - Every treatment must be reapplied after 8 days.
 - *Nit Removal* with fine-toothed comb
 - *Preventing* Reinfestation and Contagion
 - *Treat* secondary bacterial infection.

Pediculosis Corporis

Is caused by *Pediculus humanus corporis* (body louse), which resembles the head louse but is larger in size. The host invariably has very poor hygiene. The body louse lives in clothing or bedding and visits the human host only to feed. They are transmitted mainly by sharing clothes.

Clinically, nits are seen in the seams of clothing. Patient complain of generalized itching mainly on the trunk. Sites of feeding may present as pin point macules, papules, or papular urticaria. lesions are commonly scratched and can become secondarily infected.

The same topical remedies of pediculosis capitis can be used for treatment. Patient clothing and bedding must be disinfected.

Pediculosis Pubis

Pediculosis pubis is an infestation of hair-bearing regions with the pubic lice (*Phthirus pubis*). Most commonly the pubic area is affected but at times the hairy parts of the chest and axillae and the upper eyelashes but never affect the scalp

Transmission is by close physical contact such as sexual intercourse; sleeping in same bed; possibly exchange of towels. 35% of cases associated with other sexually transmitted diseases.

Clinical features:

- 1- sever itching in the pubic area, lower abdomen, upper thigh, ad sometimes in the axilla.
- 2- scratching marks.
- 3- secondary infection.
- 4- sometimes bleeding spots are seen on skin or in the underwear and may be misdiagnosed as urethral bleeding.

Diagnosis is usually clinical confirmed by finding the organism attached to hair shaft

Treatment

Shaving of the hair and application of lindane solution for 2 successive days or sulphur for 3 successive days. Treatment should be repeated after 8 days. For eyelashes infestation, application of any ointment (ex. Vaseline) to cause suffocation and death of the lice then removed by forceps.

Cutaneous Leishmaniasis (Baghdad Boil)

Leishmaniasis is a parasitic infection caused by many species of the protozoa *Leishmania*, manifested clinically as four major syndromes

- 1-Cutaneous leishmaniasis (Old and New World types).
- 2-Mucocutaneous leishmaniasis.
- 3-Diffuse cutaneous leishmaniasis.
- 4-Visceral leishmaniasis.

The old world type of cutaneous leishmaniasis is caused by:

- 1- *leishmania tropica minor* that causes the dry (urban) which heal with minimal scarring.
- 2- *leishmania tropica major* that causes the wet (rural) type which heals with ugly scarring.

(This subdivision will be our subject)

It is endemic in our country. Transmitted by the female sandfly.

The definitive host is human, while the intermediate hosts are cats, dogs, and human. Incubation period ranges from few days to few months

Clinical features: Cutaneous leishmaniasis is characterized by development of single or multiple cutaneous papules at the site of a sandfly bite, often evolving into indurated nodules with special dusky colour. These may ulcerate (in the wet type) and heal spontaneously with a depressed scar that might be ugly or distorting important organs such as the nose or the ear. It usually affects exposed areas, but may rarely affect hidden areas. Disease is self limiting healing in 9-12 months giving along lasting immunity, however recurrences may occur rarely.

Treatment

The aim is to minimize scarring and decrease the duration.

Local Therapy

- 1- Physical: Cryotherapy using liquid nitrogen.
Heating using infrared radiation.
- 2-Chemical: by intralesional injection of Na-stibogluconate, hypertonic NaCl solution or zinc sulphate.

Systemic Therapy

Used with multiple lesions, very little child or at important site like nose or angle of the eye.

- 1- Injection: Na stibogluconate (10 mg/kg/day for 10-20 days).
- 2-Oral: zinc sulphate, ketoconazole, rifampicin, INH, dapsone.