



*Citation for published version:*

Sharpe, B, Algezi, DA, Cattermole, C, Beresford, M, Bowen, R, Mitchard, J & Chalmers, AD 2017, 'A subset of high Gleason grade prostate carcinomas contain a large burden of prostate cancer syndecan-1 positive stromal cells', *Prostate*, vol. 77, no. 13, 23391, pp. 1312-1324. <https://doi.org/10.1002/pros.23391>

*DOI:*

[10.1002/pros.23391](https://doi.org/10.1002/pros.23391)

*Publication date:*

2017

*Document Version*

Peer reviewed version

[Link to publication](https://doi.org/10.1002/pros.23391)

This is the peer reviewed version of the following article: Sharpe, B., Algezi, D. A., Cattermole, C., Beresford, M., Bowen, R., Mitchard, J., & Chalmers, A. D. (2017). A subset of high Gleason grade prostate carcinomas contain a large burden of prostate cancer syndecan-1 positive stromal cells. *Prostate*, 77(13), 1312-1324, which has been published in final form at <https://doi.org/10.1002/pros.23391>. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Self-Archiving.

## University of Bath

**General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

**Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

1  
2  
3 A subset of high Gleason grade prostate carcinomas contain a large burden of prostate  
4  
5  
6 cancer syndecan-1 positive stromal cells  
7

8 Benjamin Sharpe<sup>1</sup>, Dhafer A Algezezi<sup>1,2</sup>, Claire Cattermole<sup>1</sup>, Mark Beresford<sup>3</sup>, Rebecca  
9  
10 Bowen<sup>3</sup>, John Mitchard<sup>4</sup>, Andrew D Chalmers<sup>1</sup>  
11  
12

13  
14  
15  
16 **1:** Department of Biology and Biochemistry, University of Bath, Bath, United  
17  
18 Kingdom.  
19

20  
21 **2:** Department of Medical Microbiology and Immunology, Faculty of Medicine, Thi  
22  
23 Qar University, Dhi Qar, Iraq.  
24

25 **3:** Department of Oncology, Royal United Hospital, Bath, United Kingdom.  
26

27 **4:** Department of Cellular Pathology, Royal United Hospital, Bath, United Kingdom.  
28

29  
30 Institution where work performed: Department of Biology and Biochemistry,  
31  
32 University of Bath, Bath, United Kingdom.  
33

34 Corresponding Author: Andrew Chalmers, Department of Biology and Biochemistry,  
35  
36 University of Bath, Bath, United Kingdom. Tel: 01225 385054. Fax: 01225 386779. E-mail:  
37  
38 [ac270@bath.ac.uk](mailto:ac270@bath.ac.uk).  
39

40  
41 Shortened Title: Prostate Cancer Syndecan-1 Positive stromal (PCSP) cells  
42  
43

#### 44 Disclosure Statement

45  
46  
47 The authors declare that there are no conflicts of interest to report.  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## Abstract

### Background

There is a pressing need to identify prognostic and predictive biomarkers for prostate cancer to aid treatment decisions in both early and advanced disease settings. Syndecan-1, a heparan sulfate proteoglycan, has been previously identified as a potential prognostic biomarker by multiple studies at the tissue and serum level. However, other studies have questioned its utility.

### Methods

Anti-Syndecan-1 immunohistochemistry was carried out on 157 prostate tissue samples (including cancerous, adjacent normal tissue and non-diseased prostate) from three independent cohorts of patients. A population of Syndecan-1 positive stromal cells was identified and the number and morphological parameters of these cells quantified. The identity of the Syndecan-1-positive stromal cells was assessed by multiplex immunofluorescence using a range of common cell lineage markers. Finally, the burden of Syndecan-1 positive stromal cells was tested for association with clinical parameters.

### Results

We identified a previously unreported cell type which is marked by Syndecan-1 expression and is found in the stroma of prostate tumors and adjacent normal tissue but not in non-diseased prostate. We call these cells Prostate Cancer Syndecan-1 Positive (PCSP) cells. Immunofluorescence analysis revealed that the PCSP cell population did not co-stain with markers of common prostate epithelial, stromal or immune cell populations. However, morphological analysis revealed that PCSP cells are often elongated and displayed prominent lamellipodia, suggesting they are an unidentified migratory cell population.

1  
2  
3 Analysis of clinical parameters showed that PCSP cells were found with a frequency of 20-  
4  
5 35% of all tumors evaluated, but were not present in non-diseased normal tissue.  
6

7  
8 Interestingly, a subset of primary Gleason 5 prostate tumors had a high burden of PCSP cells.  
9

## 10 Conclusions

11  
12 The current study identifies PCSP cells as a novel, potentially migratory cell type,  
13  
14 which is marked by Syndecan-1 expression and is found in the stroma of prostate  
15  
16 carcinomas, adjacent normal tissue, but not in non-diseased prostate. A subset of poor  
17  
18 prognosis high Gleason grade 5 tumors had a particularly high PCSP cell burden, suggesting  
19  
20 an association between this unidentified cell type and tumor aggressiveness.  
21  
22  
23  
24  
25  
26

## 27 Key Words:

28  
29  
30 Syndecan-1; SDC1; CD138; prostate cancer; immunohistochemistry; stromal cells  
31  
32

## 33 Introduction

34  
35  
36 Prostate cancer accounts for almost a quarter of all cancers diagnosed in European  
37  
38 men, with a large increase in incidence over the past 20 years due to increased detection  
39  
40 through PSA testing. Radical treatments such as surgery or radiotherapy have considerable  
41  
42 morbidity and there is increasing enthusiasm for a period of active surveillance in patients  
43  
44 with low volume, low-intermediate risk disease. However, we currently do not have any  
45  
46 useful biomarkers beyond Gleason grade to help inform which patients would be better  
47  
48 suited to surveillance. Likewise, in the case of metastatic prostate cancer, there is a lack of  
49  
50 useful biomarkers to help direct the choice and sequencing of drug treatments such as  
51  
52 chemotherapy, abiraterone and enzalutamide. One potential biomarker that has been  
53  
54 investigated is the heparan sulfate proteoglycan Syndecan-1.  
55  
56  
57  
58  
59  
60

1  
2  
3 The expression of Syndecan-1 has been linked to prognosis and treatment response  
4  
5 in a large range of cancer types, including hematolymphoid malignancies and solid tumors  
6  
7 (1-6), but in prostate cancer the role of Syndecan-1 in prognosis is controversial. It is  
8  
9 expressed in the basal epithelial layer of normal prostate glands, and this expression is often  
10  
11 lost in prostate carcinoma cells (7-12). However, there are reports that hormone-refractory  
12  
13 tumors (9) and tumors treated with neo-adjuvant hormone therapy (8) have increased  
14  
15 Syndecan-1 expression, suggesting a relationship between expression and androgen  
16  
17 independence. There is also evidence that expression of Syndecan-1 is associated with high  
18  
19 Gleason grade disease and/or early disease recurrence in prostate cancer patients (7-11).  
20  
21 Interestingly, an alteration in protein localization, from membranous to cytoplasmic, was  
22  
23 found to be associated with a more rapid biochemical recurrence (13). This indicates that  
24  
25 alterations in Syndecan-1 expression and localization could be prognostically important and  
26  
27 change throughout the course of the disease. Conversely, others have found that Syndecan-  
28  
29 1 expression is not associated with recurrence-free survival (9,14) or Gleason grade in  
30  
31 prostate carcinoma (9), and even concluded that lower levels of Syndecan-1 expression  
32  
33 were associated with a high Gleason grade (1). Therefore, the potential utility of Syndecan-1  
34  
35 as a biomarker for prostate cancer is far from clear.

36  
37 The function that Syndecan-1 plays in normal and cancerous tissues is also  
38  
39 complicated. Syndecan-1 protein has been linked to a wide range of cellular processes,  
40  
41 acting as an extracellular matrix receptor as well as participating in growth factor binding  
42  
43 through the heparan sulfate chains on its extracellular domain (Reviewed in (1)). The  
44  
45 extracellular domains of Syndecan-1 can also be shed through cleavage by matrix  
46  
47 metalloproteinases, creating a soluble form which may have pro-angiogenic roles in tumors,  
48  
49 for example in myeloma (1), and this shed ectodomain in itself may possess prognostic  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 utility (12). There is also evidence that Syndecan-1 signaling is involved in maintaining the  
4  
5 tumor-initiating cell/cancer stem cell population in breast carcinoma (15,16) and prostate  
6  
7 carcinoma (17) and can regulate the expression of several miRNAs that suppress expression  
8  
9 of stem cell markers such as SOX-2, OCT4 and Nanog (11). Suppression of Syndecan-1 gene  
10  
11 expression also reduces amplification of the putative tumor-initiating cell population  
12  
13 following treatment with docetaxel (17), suggesting that it could exert effects of relevance  
14  
15 to prostate cancer patients undergoing therapy. Taken together, there is accumulating  
16  
17 evidence that Syndecan-1 is an important multifunctional molecule that shows altered  
18  
19 expression in prostate cancer, but its potential role as a biomarker for prostate cancer  
20  
21 remains controversial.  
22  
23  
24  
25

26  
27 Given the interesting functional roles played by Syndecan-1 and the conflicting  
28  
29 evidence surrounding its possible utility as a biomarker, we investigated Syndecan-1  
30  
31 expression in three independent cohorts of prostate carcinoma samples. We identified a  
32  
33 previously unreported cell type which is marked by Syndecan-1 expression and is found  
34  
35 scattered in the stroma of a third of prostate carcinomas and in their adjacent normal tissue,  
36  
37 but not in normal tissue. We call these cells Prostate Cancer Syndecan-1 Positive (PCSP) cells.  
38  
39 Common markers for immune cells, cancer epithelial cells and stromal cells failed to identify  
40  
41 the cell type, but further analysis showed that PCSP cells are elongated and have prominent  
42  
43 lamellipodia, suggesting they are a migratory cell type. Finally, comparison of the  
44  
45 relationship between PCSP cell burden and clinical features showed that a subset of poor  
46  
47 prognosis high grade tumors have a particularly high burden of these cells.  
48  
49  
50  
51  
52

## 53 Materials and Methods

### 56 Patients

1  
2  
3 The study was conducted following local ethics approval (REC reference:  
4  
5 13/WS/0153; IRAS project ID: 112241). Data was obtained from three separate cohorts of  
6  
7 patients with associated formalin-fixed, paraffin-embedded (FFPE) prostate material  
8  
9 available. The total number of cancer cases in the study was 157. The Bath cohort was used  
10  
11 for analysis of cell types by co-staining, and consisted of 6 prostate cancer patients with  
12  
13 confirmed Syndecan-1+ stromal cells as assessed by anti-Syndecan-1 IHC. In addition, one  
14  
15 block of adjacent normal prostate tissue was used as a positive control for Syndecan-1  
16  
17 staining. These samples were obtained retrospectively from patients treated for prostate  
18  
19 cancer at the Royal United Hospital, Bath, United Kingdom, between 1997 and 2008. The  
20  
21 other two cohorts, consisting of 151 cancer cases in total, were used for anti-Syndecan-1  
22  
23 immunohistochemistry and were obtained from US Biomax (Rockville, MD, USA). The first  
24  
25 cohort (PR1921) consisted of 80 cases of prostate adenocarcinoma with 8 cases of adjacent  
26  
27 normal prostate tissue and 8 cases of histologically normal prostate tissue, with duplicate  
28  
29 1mm diameter cores per cancer case for a total of 196 cores. The second cohort (PR803b)  
30  
31 consisted of 71 cases of prostate adenocarcinoma, 2 cases of prostate leiomyosarcoma, 1  
32  
33 case of benign prostate hyperplasia and 6 cores of histologically normal prostate, with a  
34  
35 single 1.5mm core per case. Each cohort was contained in a single tissue microarray block. A  
36  
37 summary of the patient characteristics of tissue microarray cohorts is given in Table I.  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

### Immunohistochemistry

For immunohistochemistry, 5µm thick sections of FFPE prostate tissue or tissue microarrays were used. Cohort 1 and cohort 2 were stained in separate experiments. Briefly, sections were baked for 30 minutes at 60°C before being deparaffinized in Histoclear (National Diagnostics, Hesse, UK) and rehydrated through a graded ethanol series of decreasing ethanol concentration. Sections were washed with PBS-tween (0.05%), permeabilized with 0.5% triton X-100 in PBS, subjected to heat-induced epitope retrieval in 10mM Citrate buffer with 0.05% Tween 20 at pH 6.0 for 30 minutes at 90°C, and allowed to cool to room temperature for 20 minutes. Endogenous peroxidases were blocked with Dako Peroxidase Block, following the manufacturer's instructions. After blocking for 30 minutes in 10% normal goat serum and 0.5% BSA in PBS, samples were incubated with primary antibodies (Table II) diluted in Dako Antibody Diluent (Dako, Ely, UK) overnight at 4°C. The next day, samples were washed and bound antibodies were visualized using the EnVision+ Kit (K400611-2 and K401011-2, Dako, Ely, UK) following the manufacturer's instructions with DAB exposure for 5 minutes. Nuclei were counterstained with Gill's Hematoxylin (H-3401, Vector Laboratories, Peterborough, UK) and mounted in DPX (Sigma-Aldrich, Dorset, UK). Stained tissues were viewed under a Nikon Eclipse E800 microscope with brightfield illumination and photographed with a Nikon Digital Sight DS-U1 CCD Digital Camera. Negative controls were IHC-stained sections of prostate carcinoma tissue with no primary antibody added, and showed no positive staining (data not shown).

#### Anti-Syndecan-1 Antibodies

Three distinct Syndecan-1 antibodies were used (Table II). The three antibodies were selected as they have been previously shown to successfully detect Syndecan-1. The first mouse monoclonal anti-Syndecan-1 antibody, (Novus Biologicals: clone B-A38: Cat NB100-



1  
2  
3 64980), later referred to as mo anti-SDC1 #1, has been shown to robustly mark a CD45-  
4  
5 expressing plasma cell population by flow cytometry (18), a cell population which expresses  
6  
7 Syndecan-1 (19). The second mouse monoclonal (Dako; clone MI15: Cat M7228) has been  
8  
9 shown to recognize the ectodomain of Syndecan-1 (20). Antibody blocking experiments  
10  
11 show that it partially shares the epitope of B-B4, which occurs at residues 90-93 of the  
12  
13 mature Syndecan-1 protein (21). The rabbit polyclonal anti-Syndecan-1 antibody (Santa Cruz  
14  
15 Biotechnology: SC-5632) was generated by immunization with amino acids 82-256 of human  
16  
17 Syndecan-1, corresponding to the human Syndecan-1 ectodomain, and successfully  
18  
19 immunoprecipitated the ectodomain (22), showing that the antibody can detect the cleaved  
20  
21 ectodomain of the protein. The rabbit antibody is also known to detect Syndecan-1 positive  
22  
23 plasma cell populations by immunohistochemistry (23).  
24  
25  
26  
27

### 28 **Scoring**

29  
30  
31 For assessment of IHC staining on the two tissue microarrays, whole cores were  
32  
33 examined under a 10x objective to determine the presence of Syndecan-1 staining in  
34  
35 prostate epithelium. Cores were scored as either positive or negative for Syndecan-1  
36  
37 epithelial reactivity based on the observed staining patterns across the core.  
38  
39

40  
41 For quantification of PCSP cell burden, fields of view of all PCSP cells in every core  
42  
43 were manually acquired under a 20x objective. Counting and measuring of the resulting  
44  
45 images was performed using ImageJ v2.0.0. Each PCSP cell was identified manually and the  
46  
47 following size and shape descriptors were measured: area, circularity, aspect ratio,  
48  
49 roundness and solidity. Mean values were taken for each measure within each cohort as a  
50  
51 whole, and for each patient.  
52  
53  
54

### 55 **Data Analysis**

1  
2  
3 To analyze the distribution of Syndecan-1+ cell burden amongst different patient  
4  
5 groups in both TMA cohorts, we classified patients into Syndecan-1 stromal cell positive and  
6  
7 negative categories based on whether Syndecan-1+ stromal cells were present or absent in  
8  
9 the core. In the case of PR1921, where duplicate cores were scored, patients were classified  
10  
11 as positive if at least one core contained Syndecan-1+ stromal cells. The distributions of  
12  
13 Syndecan-1+ and Syndecan-1- patients amongst different levels of clinical variables were  
14  
15 tested with the chi-squared test performed in R version 3.2.0 with default parameters.  
16  
17 Clinical variables assessed were: age; stage; Gleason grade; T, N and M stages; Presence of  
18  
19 Syndecan-1 staining in the glandular compartment was also scored as positive or negative,  
20  
21 and tested for association as with clinical variables. A p-value of less than 0.05 was  
22  
23 considered statistically significant. Graphs were generated in R version 3.2.0.  
24  
25  
26  
27

### 28 29 **Immunofluorescence**

30  
31 For immunofluorescence co-staining of Syndecan-1 with other markers, double-  
32  
33 labelling was performed on FFPE prostatectomies, biopsies and TURP chips with confirmed  
34  
35 PCSP cells as determined by prior IHC experiments. 5um thick sections were deparaffinized,  
36  
37 rehydrated, permeablized and subjected to antigen retrieval as for immunohistochemistry.  
38  
39 Omitting the peroxidase blocking step, samples were blocked with the same blocking buffer  
40  
41 and incubated with a mixture of two primary antibodies raised in different species,  
42  
43 overnight at 4°C, as before. Cell type markers and either the rabbit polyclonal or the first  
44  
45 mouse monoclonal (mo anti-SDC1 #1) anti-Syndecan-1 antibodies were chosen, depending  
46  
47 on the host in which the marker antibody was raised. The antibodies used in this study are  
48  
49 described in Table II. The following day, samples were washed and incubated in a secondary  
50  
51 antibody cocktail containing goat anti-mouse alexafluor-488 (A-11001, Invitrogen) and goat  
52  
53 anti-rabbit alexafluor-568 (A-11008, Invitrogen), each at 1:200 dilutions in Dako Antibody  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 Diluent, for 30 minutes. The specimens were then incubated with a solution of 0.1% (w/v)  
4  
5 Sudan Black B (199664, Sigma-Aldrich, Dorset, UK) in 70% ethanol for 10 minutes to quench  
6  
7 endogenous tissue autofluorescence, as adapted from (24). Mounting was achieved in  
8  
9 Mowiol-4-88 (Millipore) with 600nM DAPI (D9542, Sigma-Aldrich) added. Slides were  
10  
11 allowed to set overnight in the dark at room temperature before imaging on a ZEISS LSM510  
12  
13 META confocal laser scanning microscope. Negative controls were IF-stained sections of  
14  
15 prostate carcinoma tissue with no primary antibody added, and never showed positive  
16  
17 staining (data not shown).  
18  
19  
20  
21

22 To score Syndecan-1+ stromal cells for their colocalization with test markers, images  
23  
24 from each experimental run were pooled, and each Syndecan-1+ cell was manually scored  
25  
26 for positivity of the co-stained marker. The percentage of marker-positive cells was  
27  
28 calculated as the percentage of Syndecan-1+/Marker+ cells relative to the total number of  
29  
30 Syndecan-1+ cells in an experimental run.  
31  
32  
33

## 34 Results

### 35 36 37 **A population of Syndecan-1 positive cells was found in the stromal compartment** 38 39 **of a subset of prostate tumors** 40 41

42 Syndecan-1 expression was analyzed using IHC and IF in samples from three  
43  
44 independent cohorts of patients (See Materials and Methods). Expression was found in the  
45  
46 cell membrane and cytoplasm of the prostate epithelium basal, but not luminal, cell layer in  
47  
48 histologically normal tissue (Figure 1A, Left Panel). In prostate carcinoma tissues there was  
49  
50 either a loss of epithelial staining or a widespread expression of the proteoglycan  
51  
52 throughout the cytoplasm and/or cell membranes of prostate tumor epithelium (Figure 1A,  
53  
54 Middle and right Panel). These patterns are consistent with previous studies  
55  
56  
57  
58  
59  
60

1  
2  
3 (3,7,10,12,13,25) and provide positive (basal prostate epithelium) and negative (luminal  
4  
5 prostate epithelium) control cells for the IHC staining. Interestingly, in addition to these  
6  
7 staining patterns, a population of scattered Syndecan-1 positive cells was found in the  
8  
9 stroma in a subset of cancer cases (Figure 1A). These scattered cells were present in a  
10  
11 subset of tumor samples from each of the three cohorts, either proximal or distal to an  
12  
13 epithelial region. A range of examples, including samples from all three cohorts, are shown  
14  
15 in Figure 1B. The staining was found to be localized to the membranes and/or cytoplasmic  
16  
17 regions of the Syndecan-1 positive cells, and the cells often appeared in close proximity to  
18  
19 areas of apparent inflammation, characterized by the presence of stromal cells with round,  
20  
21 compact and hyperchromatic properties suggestive of an immune cell phenotype (Figure 1B,  
22  
23 Adjacent to stained cells in the inserts and near arrowheads. The bottom left panel insert  
24  
25 shows a clear example).  
26  
27  
28  
29  
30

31 To confirm that the staining was due to the presence of Syndecan-1, and not cross  
32  
33 reactivity from the mouse monoclonal antibody (Clone B-A38), two additional anti-  
34  
35 Syndecan-1 antibodies were used. A rabbit polyclonal antibody and a second mouse  
36  
37 monoclonal (Clone MI15) were used. Staining of sequential sections of normal prostate and  
38  
39 tonsil by IHC showed that all three antibodies stained the same basal, but not luminal,  
40  
41 epithelial cell populations (Figure 2A and B). This specific pattern of expression provides  
42  
43 positive and negative controls for the three antibodies and, combined with validation of the  
44  
45 antibodies carried out in previous publications (see materials and methods), argues that  
46  
47 they can be used to reliably identify Syndecan-1 positive cells. The two additional anti-  
48  
49 Syndecan-1 antibodies (rabbit polyclonal and second mouse monoclonal), like the original  
50  
51 mouse monoclonal antibody, also detected scattered Syndecan-1 positive cells in the  
52  
53 stroma of prostate tumors (Figure 2C).  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 A final experiment to validate the anti-syndecan-1 antibodies was carried out using  
4  
5 IF staining of normal and prostate carcinoma tissues. This made it possible to establish if the  
6  
7 different antibodies were staining the same populations at a single cell level. IF staining  
8  
9 using the mouse monoclonal #1 and the rabbit polyclonal antibody showed colocalisation of  
10  
11 Syndecan-1 staining in individual basal, but not luminal, epithelial cells of adjacent normal  
12  
13 prostate (Figure 3, upper panels). Importantly, colocalization of Syndecan-1 staining was  
14  
15 also seen in the scattered tumor stromal cells (Figure 3, lower panels arrows and inserts),  
16  
17 confirming that the two antibodies stain the same population of scattered stromal cells.  
18  
19  
20  
21

22 In summary, the use of three independent antibodies showed that a subset of  
23  
24 prostate cancer samples, from three cohorts of patients, contained a population of  
25  
26 scattered Syndecan-1 positive cells, which to our knowledge have not been previously  
27  
28 described. We call these cells Prostate Cancer Syndecan-1 Positive (PCSP) cells.  
29  
30  
31  
32

### 33 **PCSP cells do not express common epithelial, mesenchymal or immune cell** 34 35 **markers**

36  
37  
38 To begin to characterize the PCSP cell population, we used double-IF labelling with  
39  
40 Syndecan-1 and a range of markers commonly used to distinguish key cell types found in  
41  
42 prostate tumor samples from the Bath cohort. Given that Syndecan-1 is expressed in normal  
43  
44 and neoplastic plasma cells (19), we began by assessing the expression of hematolymphoid  
45  
46 lineage markers in PCSP cells. Fluorescent co-stains with Syndecan-1 and a panel of  
47  
48 antibodies raised against immune cell lineage markers: CD45, which is expressed in the  
49  
50 majority of hematolymphoid lineages with the exception of erythrocytes and some  
51  
52 macrophages, and known to be expressed in Syndecan-1+ bone marrow plasma cells (18);  
53  
54 CD19, which is expressed on immature and memory B-cells (26); CD27, which is expressed in  
55  
56  
57  
58  
59  
60

1  
2  
3 plasma cells, memory B-cells (26) and T-cells (27); and CD73, an ecto-5'-nucleotidase  
4  
5 expressed on subsets of T and B lymphocytes (28), as well mesenchymal stem cells which  
6  
7 have been previously shown to infiltrate prostate tumors (29). To confirm the antibodies for  
8  
9 these markers were performing as expected they were used to stain positive control tonsil  
10  
11 or placenta tissues (Supplementary Figure 1). In each case the staining was as expected.  
12  
13

14  
15 In prostate tumors, the majority of PCSP cells did not co-stain for any of the three  
16  
17 immune cell markers or the mesenchymal stem cell marker (Figure 4A+B). Interestingly,  
18  
19 despite the lack of colocalization, PCSP cells were often found in close proximity to CD45-  
20  
21 positive immune cells (Figure 4A: See inserts in the top panel). This was consistent with IHC  
22  
23 data showing that Syndecan-1 positive cells were found in close proximity to cells whose  
24  
25 nuclear morphology was consistent with an immune cell phenotype (Figure 1B, Adjacent to  
26  
27 stained cells in the inserts and near arrowheads. The bottom left panel insert shows a clear  
28  
29 example).  
30  
31  
32

33  
34 Given that Syndecan-1 is expressed in the basal cells of normal prostate, another  
35  
36 possible origin for the PCSP population was prostate epithelial cells, so co-staining was  
37  
38 carried out for Syndecan-1 and three markers of epithelial cell fate: pan-Cytokeratin, which  
39  
40 stains multiple human cytokeratins expressed in epithelial tissues including both basal and  
41  
42 luminal prostate epithelial cell layers (Figure 5A); E-Cadherin, which is normally expressed  
43  
44 throughout the normal prostate epithelium and in most prostate tumors (30); and Prostate-  
45  
46 Specific Antigen (PSA), a secretory marker of luminal prostate epithelium, also expressed in  
47  
48 prostate tumors. Quantification of the IF images showed none of these markers had  
49  
50 significant co-staining with Syndecan-1 in PCSP cells (Figure 5B). There was also no apparent  
51  
52 overlap in the IHC staining of the stromal cells in consecutive tissue sections for Syndecan-1  
53  
54 and the basal cell markers p63 and Cytokeratin-5 (Figure 5C, lower panels). This was clearly  
55  
56  
57  
58  
59  
60

1  
2  
3 distinct from the situation in the basal cells of adjacent normal tissue, where the markers do  
4  
5 overlap with Syndecan-1 staining (Figure 5C, upper panels). Thus, despite the overlap in  
6  
7 normal epithelium, PCSP cells did not have staining patterns consistent with them being a  
8  
9 recognized prostate epithelial cell population, whether basal or luminal in origin.

10  
11  
12 To examine if the PCSP cells were stromal, mesenchymal or had undergone an  
13  
14 epithelial-mesenchymal transition and lost expression of epithelial markers, a panel of  
15  
16 stains for mesenchymal and stromal cell types was examined. The mesenchymal cell  
17  
18 markers Vimentin and N-Cadherin did not show significant co-staining with PCSP cells  
19  
20 (Figure 6A+B). Finally, the neural marker S100 and the endothelial marker CD31 were  
21  
22 examined (Figure 6A), neither of which showed significant overlap (Figure 6B). Therefore,  
23  
24 the marker analysis carried out did not allow the allocation of PCSP cells to an established  
25  
26 cell population, and they remained an undefined cell type.  
27  
28  
29  
30  
31  
32

### 33 **PCSP cells are elongated and have prominent lamellipodia-like structures**

34  
35  
36 In order to characterize this unidentified cell type further, the shape of PCSP stromal  
37  
38 cells was analyzed using IHC stained tissue microarrays from two independent cohorts of  
39  
40 prostate cancer patients. The circularity and aspect ratios of the PCSP cells were quantified  
41  
42 and on average PCSP cells were found to be elongated rather than rounded in shape (Table  
43  
44 III). When morphological data from individual cells of both TMA cohorts were pooled, a tail  
45  
46 of elongated cells with high aspect ratio and low circularity was apparent (Figure 7A). To  
47  
48 further characterize this subpopulation of elongated cells, confocal laser scanning  
49  
50 microscopy was used to carry out high resolution three dimensional imaging of the PCSP  
51  
52 cells. They were found to have striking lamellipodia-like protrusions (Figure 7B, arrows) and  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 polarized cell bodies and nuclei (Figure 7B, arrowheads). The elongated shape and  
4  
5 prominent protrusions support the hypothesis that PCSP cells may be a migratory cell type.  
6  
7

### 8 9 10 **A subset of Gleason grade 5 tumors had a particularly high PCSP cell burden**

11  
12 Using the clinical metadata available for both prostate tissue microarray cohorts, we  
13  
14 wished to determine if there was a relationship between PCSP cell burden and  
15  
16 clinicopathological features. These included age, Gleason grade and TNM stages. In the first  
17  
18 cohort, cores of histologically normal tissue adjacent to prostate cancer were also included,  
19  
20 as well as non-diseased normal prostate control cores. Patient characteristics for both  
21  
22 cohorts are summarized in Table I, and have similar patient distributions for the clinical  
23  
24 variables.  
25  
26  
27

28  
29 PCSP cells were found in 28 of 80 cases (35%) of patients with prostate cancer in  
30  
31 cohort 1, but they were not observed in any of the 8 control cases of non-diseased patient  
32  
33 tissue (Figure 8A). PCSP cells were found in 4 of 8 cases of normal patient tissue that was  
34  
35 adjacent to carcinoma (Figure 8A). In cohort 2, PCSP cells were found in 15 of 71 prostate  
36  
37 cancer cases (21.1%). In both cohorts, a subset of Gleason grade five tumors had a  
38  
39 particularly high burden of PCSP cells (Figure 8B+C). Among cohort 2, but not cohort 1, there  
40  
41 was also a statistically significant association between PCSP cell burden and higher primary  
42  
43 Gleason grade ( $p < 0.05$ ). In summary, analysis of PCSPs in the tissue microarrays showed  
44  
45 that PCSP cells were not observed in normal prostate tissue, but were found in adjacent  
46  
47 normal and in roughly 1/3 of prostate carcinoma tissues, with a subset of higher grade  
48  
49 tumors having a particularly high burden.  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



## Discussion

This study identifies PCSP cells as a previously unreported Syndecan-1-positive cell population found in the stroma of prostate adenocarcinoma, in adjacent normal tissue, but not in non-diseased prostate. These PCSP cells do not express lineage markers of common epithelial, stromal or immune cell lineages, but do display elongated morphologies with lamellipodia-like structures consistent with a migrating cell type. PCSP cells appear in high burdens in a subset of primary Gleason grade 5 poor prognosis tumors.

The existence of PCSP cells was confirmed using three separate antibodies recognizing Syndecan-1; in both immunohistochemistry and immunofluorescence, using samples from three different patient cohorts. However, several previous studies have specifically reported the absence of stromal Syndecan-1 staining in prostate tumors (7,10,12). It is possible that the focal nature of this cell type – and their presence only in a subset of tumors - has led to them being previously overlooked.

Immune cells are commonly found in prostate tumors and Syndecan-1 is a marker of plasma cell differentiation, being expressed at the later stages of the transition from plasmablast into a mature plasma cell (31). However, we were unable to identify an immune cell marker profile in PCSP cells. The fact that plasma cells tend to express both Syndecan-1 and the pan-leukocyte marker CD45 (18), whilst PCSP cells are CD45-, would suggest that PCSP cells are either an atypical plasma cell or a non-plasma cell population. Tumor-associated macrophages - a component commonly associated with inflammation in tumors – also express CD45, being a leukocyte population, and that makes it unlikely that PCSP cells represent a subset of that population. However, given the enormous diversity in the

1  
2  
3 immune cell repertoire, it is still possible that PCSP cells represent an immune cell  
4  
5 population.  
6

7  
8 Co-staining with epithelial markers again failed to mark a significant portion of the  
9  
10 PCSP cell population, despite the overlap seen in adjacent normal epithelium. There remains  
11  
12 the possibility that PCSP cells could have undergone dedifferentiation, as they do not  
13  
14 possess an epithelial identity consistent with tumor cells. Epithelial-Mesenchymal Transition  
15  
16 (EMT) is an unlikely explanation for this observation, as PCSP cells also lacked mesenchymal  
17  
18 markers such as Vimentin and N-Cadherin (Figure 5A+B). Another possibility for the lack of  
19  
20 differentiation markers is a stem-like status, either derived from a resident prostate cell  
21  
22 population or from a mobilized stem cell niche such as Mesenchymal Stem Cells (32) or  
23  
24 cancer stem cells, although co-staining with a Mesenchymal Stem Cell marker (CD73) did  
25  
26 not show any colocalisation with PCSP cells. There is evidence that Syndecan-1 signaling is  
27  
28 involved in maintaining the tumor-initiating cell/cancer stem cell population in the prostate  
29  
30 carcinoma cell line PC-3, where its expression marks a population of CD133+/CD44+ stem-  
31  
32 like holoclones (17), so a migratory dedifferentiated/stem like epithelial cell identity  
33  
34 remains a possibility. It is also possible that Syndecan-1 might stain more than one  
35  
36 population of cells in the stroma that overlap with different markers. We feel this is a less  
37  
38 likely than a single population of PCSP cells, but a key question for future work is to identify  
39  
40 the markers expressed by these cells to establish their identity and whether they represent  
41  
42 more than one population of cells.  
43  
44  
45  
46  
47  
48

49  
50 Analysis of PCSP morphology suggests a migratory cell type, something that is  
51  
52 supported by the finding that histologically normal tissue adjacent to a carcinoma can  
53  
54 harbor PCSP cells. A migratory PCSP cell population could be either recruited from other  
55  
56 tissues, consistent with an immune cell type; a hypothesis which is supported by the close  
57  
58  
59  
60

1  
2  
3 proximity of other immune cells to the PCSP cells. However, PCSP cells might also be  
4  
5 migrating out from the prostate tumor and recruiting or being detected by immune cells,  
6  
7 which would be more supportive of a dedifferentiated/stem like epithelial population.  
8  
9

10 In addition to their characterization, it is important to determine the clinical  
11  
12 relevance of PCSP cells. We observed that there is a trend towards increased burden in the  
13  
14 primary Gleason grade 5 tumors, suggesting that PCSP burden might be related to outcome  
15  
16 given that assignment of primary Gleason grade 5 is a poor prognostic factor. While cohort 2  
17  
18 showed a significant association between primary Gleason grade and the presence of a  
19  
20 PCSP cell burden ( $p < 0.05$ ), cohort 1 did not reach significance ( $p = 0.706$ ). This could be due  
21  
22 to differences in sampling strategy between the two cohorts because of different core  
23  
24 numbers and diameters, with cohort 1 having duplicate 1mm cores and cohort 2 having  
25  
26 single 1.5mm cores. This could lead to differences in successful detection of Syndecan-1+  
27  
28 cells in tumors given their scattered nature. Further investigation is warranted to determine  
29  
30 the relationship between PCSP burden and grade, and prognostic utility of PCSP cells in  
31  
32 prostate carcinoma, as well as any potential for predicting which cancers may respond  
33  
34 better to upfront systemic cytotoxic or hormone treatments, and which may benefit from  
35  
36 active surveillance.  
37  
38  
39  
40  
41  
42  
43

## 44 Conclusions

45  
46  
47 This report is, we believe, the first to identify PCSP cells, an apparently migratory  
48  
49 Syndecan-1+ cell population found in the stroma of prostate tumors and adjacent normal  
50  
51 tissue, but not in non-diseased prostate. In addition, we demonstrate an increased burden  
52  
53 of this cell type in primary poor prognosis Gleason pattern 5 tumors. These findings provide  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 a foundation for further characterization of PCSP cells, their clinical relevance and their roles  
4  
5 in prostate tumors and the adjacent normal tissue.  
6  
7

## 8 9 Acknowledgements

10  
11 The authors would like to thank Professor Robert Kelsh and the Bioimaging Suite at  
12  
13 the University of Bath for providing use of imaging facilities and Doctor Gordon Taylor for  
14  
15 statistical advice. The authors are grateful to the Annett Trust for research project grant  
16  
17 funding and the Iraq government for PhD scholarship funding of D.A. Algezei. They are  
18  
19 members of the Cancer Research at Bath network, which provided seedcorn grant funding  
20  
21 for this project. Lastly, they are grateful to the Department of Cellular Pathology at the  
22  
23 Royal United Hospital, Bath, for specimen handling and curation.  
24  
25  
26  
27  
28

## 29 30 References

- 31  
32  
33 1. Gharbaran R. Advances in the molecular functions of syndecan-1 (SDC1/CD138) in  
34 the pathogenesis of malignancies. *Critical Reviews in Oncology Hematology*  
35 2015;94(1):1-17.
- 36 2. Nguyen TL, Grizzle WE, Zhang K, Hameed O, Siegal GP, Wei S. Syndecan-1  
37 Overexpression Is Associated With Nonluminal Subtypes and Poor Prognosis in  
38 Advanced Breast Cancer. *American Journal of Clinical Pathology* 2013;140(4):468-  
39 474.
- 40 3. Zellweger T, Ninck CH, Mirlacher M, Annefeld M, Glass AG, Gasser TC, Mihatsch MJ,  
41 Gelmann EP, Bubendorf L. Tissue microarray analysis reveals prognostic significance  
42 of syndecan-1 expression in pro-state cancer. *Prostate* 2003;55(1):20-29.
- 43 4. Juuti A, Nordling S, Lundin J, Louhimo J, Haglund C. Syndecan-1 expression - A novel  
44 prognostic marker in pancreatic cancer. *Oncology* 2005;68(2-3):97-106.
- 45 5. Lee SH, Choi EJ, Kim MS, Park JW, Lee YS, Kim SY, Kang CS. Prognostic significance of  
46 syndecan-1 expression in squamous cell carcinoma of the tonsil. *International*  
47 *Journal of Clinical Oncology* 2014;19(2):247-253.
- 48 6. Ramani VC, Sanderson RD. Chemotherapy stimulates syndecan-1 shedding: A  
49 potentially negative effect of treatment that may promote tumor relapse. *Matrix*  
50 *Biology* 2014;35:215-222.
- 51 7. Chen D, Adenekan B, Chen L, Vaughan ED, Gerald W, Feng ZD, Knudsen BS.  
52 Syndecan-1 expression in locally invasive and metastatic prostate cancer. *Urology*  
53 2004;63(2):402-407.  
54  
55  
56  
57  
58  
59  
60

- 1
  - 2
  - 3
  - 4
  - 5
  - 6
  - 7
  - 8
  - 9
  - 10
  - 11
  - 12
  - 13
  - 14
  - 15
  - 16
  - 17
  - 18
  - 19
  - 20
  - 21
  - 22
  - 23
  - 24
  - 25
  - 26
  - 27
  - 28
  - 29
  - 30
  - 31
  - 32
  - 33
  - 34
  - 35
  - 36
  - 37
  - 38
  - 39
  - 40
  - 41
  - 42
  - 43
  - 44
  - 45
  - 46
  - 47
  - 48
  - 49
  - 50
  - 51
  - 52
  - 53
  - 54
  - 55
  - 56
  - 57
  - 58
  - 59
  - 60
8. Shimada K, Nakamura M, De Velasco MA, Tanaka M, Ouji Y, Konishi N. Syndecan-1, a new target molecule involved in progression of androgen-independent prostate cancer. *Cancer Science* 2009;100(7):1248-1254.
9. Brimo F, Vollmer RT, Friszt M, Corcos J, Bismar TA. Syndecan-1 expression in prostate cancer and its value as biomarker for disease progression. *Bju International* 2010;106(3):418-423.
10. Shariat SF, Svatek RS, Kabbani W, Walz J, Lotan Y, Karakiewicz PI, Roehrborn CG. Prognostic value of syndecan-1 expression in patients treated with radical prostatectomy. *Bju International* 2008;101(2):232-237.
11. Fujii T, Shimada K, Tatsumi Y, Fujimoto K, Konishi N. Syndecan-1 responsive microRNA-126 and 149 regulate cell proliferation in prostate cancer. *Biochemical and biophysical research communications* 2015;456(1):183-189.
12. Szarvas T, Reis H, vom Dorp F, Tschirdewahn S, Niedworok C, Nyirady P, Schmid KW, Rubben H, Kovalszky I. Soluble syndecan-1 (SDC1) serum level as an independent pre-operative predictor of cancer-specific survival in prostate cancer. *Prostate* 2016;76(11):977-985.
13. Ledezma R, Cifuentes F, Gallegos I, Fulla J, Ossandon E, Castellon EA, Contreras HR. Altered expression patterns of syndecan-1 and -2 predict biochemical recurrence in prostate cancer. *Asian Journal of Andrology* 2011;13(3).
14. Purushothaman A, Uyama T, Kobayashi F, Yamada S, Sugahara K, Rapraeger AC, Sanderson RD. Heparanase-enhanced shedding of syndecan-1 by myeloma cells promotes endothelial invasion and angiogenesis. *Blood* 2010;115(12):2449-2457.
15. Alexander CM, Reichsman F, Hinkes MT, Lincecum J, Becker KA, Cumberledge S, Bernfield M. Syndecan-1 is required for Wnt-1-induced mammary tumorigenesis in mice. *Nature Genetics* 2000;25(3):329-332.
16. Liu BY, McDermott SP, Khwaja SS, Alexander CM. The transforming activity of Wnt effectors correlates with their ability to induce the accumulation of mammary progenitor cells. *Proceedings of the National Academy of Sciences of the United States of America* 2004;101(12):4158-4163.
17. Shimada K, Anai S, Fujii T, Tanaka N, Fujimoto K, Konishi N. Syndecan-1 (CD138) contributes to prostate cancer progression by stabilizing tumour-initiating cells. *Journal of Pathology* 2013;231(4):495-504.
18. Maiga RI, Lemieux J, Roy A, Simard C, Neron S. Flow Cytometry Assessment of In Vitro Generated CD138(+) Human Plasma Cells. *Biomed Research International* 2014.
19. Chilosi M, Adami F, Lestani M, Montagna L, Cimarosto L, Semenzato G, Pizzolo G, Menestrina F. CD138/syndecan-1: A useful immunohistochemical marker of normal and neoplastic plasma cells on routine trephine bone marrow biopsies. *Modern Pathology* 1999;12(12):1101-1106.
20. Gattei V, Godeas C, Degan M, Rossi FM, Aldinucci D, Pinto A. Characterization of anti-CD138 monoclonal antibodies as tools for investigating the molecular polymorphism of syndecan-1 in human lymphoma cells. *Br J Haematol* 1999;104(1):152-162.
21. Dore JM, Morard F, Vita N, Wijdenes J. Identification and location on syndecan-1 core protein of the epitopes of B-B2 and B-B4 monoclonal antibodies. *FEBS Lett* 1998;426(1):67-70.
22. Schmidt A, Echtermeyer F, Alozie A, Brands K, Buddecke E. Plasmin- and thrombin-accelerated shedding of syndecan-4 ectodomain generates cleavage sites at

- 1  
2  
3 Lys(114)-Arg(115) and Lys(129)-Val(130) bonds. *Journal of Biological Chemistry*  
4 2005;280(41):34441-34446.
- 5 23. Huard B, McKee T, Bosshard C, Durual S, Matthes T, Myit S, Donze O, Frossard C,  
6 Chizzolini C, Favre C, Zubler R, Guyot JP, Schneider P, Roosnek E. APRIL secreted by  
7 neutrophils binds to heparan sulfate proteoglycans to create plasma cell niches in  
8 human mucosa. *Journal of Clinical Investigation* 2008;118(8):2887-2895.
- 9 24. Viegas MS, Martins TC, Seco F, do Carmo A. An improved and cost-effective  
10 methodology for the reduction of autofluorescence in direct immunofluorescence  
11 studies on formalin-fixed paraffin-embedded tissues. *European Journal of*  
12 *Histochemistry* 2007;51(1):59-66.
- 13 25. Contreras HR, Ledezma RA, Vergara J, Cifuentes F, Barra C, Cabello P, Gallegos I,  
14 Morales B, Huidobro C, Castellon EA. The expression of syndecan-1 and-2 is  
15 associated with Gleason score and epithelial-mesenchymal transition markers, E-  
16 cadherin and beta-catenin, in prostate cancer. *Urologic Oncology-Seminars and*  
17 *Original Investigations* 2010;28(5).
- 18 26. Mavropoulos A, Simopoulou T, Varna A, Liaskos C, Katsiari CG, Bogdanos DP, Sakkas  
19 LI. Breg Cells Are Numerically Decreased and Functionally Impaired in Patients With  
20 Systemic Sclerosis. *Arthritis & Rheumatology* 2016;68(2):494-504.
- 21 27. Hendriks J, Gravestien LA, Tesselaar K, van Lier RAW, Schumacher TNM, Borst J.  
22 CD27 is required for generation and long-term maintenance of T cell immunity.  
23 *Nature Immunology* 2000;1(5):433-440.
- 24 28. Resta R, Yamashita Y, Thompson LF. Ecto-enzyme and signaling functions of  
25 lymphocyte CD73. *Immunological Reviews* 1998;161:95-109.
- 26 29. Brennen WN, Chen S, Denmeade SR, Isaacs JT. Quantification of Mesenchymal Stem  
27 Cells (MSCs) at Sites of Human Prostate Cancer. *Oncotarget* 2013;4(1):106-117.
- 28 30. Rubin MA, Mucci NR, Figurski J, Fecko A, Pienta KJ, Day ML. E-cadherin expression in  
29 prostate cancer: A broad survey using high-density tissue microarray technology.  
30 *Human Pathology* 2001;32(7):690-697.
- 31 31. De Vos J, Hose D, Reme T, Tarte K, Moreaux J, Mahtouk K, Jourdan M, Goldschmidt H,  
32 Rossi JF, Cremer FW, Klein B. Microarray-based understanding of normal and  
33 malignant plasma cells. *Immunological Reviews* 2006;210:86-104.
- 34 32. Cheng J, Yang K, Zhang Q, Yu Y, Meng Q, Mo N, Zhou Y, Yi X, Ma C, Lei A, Liu Y. The  
35 role of mesenchymal stem cells in promoting the transformation of androgen-  
36 dependent human prostate cancer cells into androgen-independent manner.  
37 *Scientific Reports* 2016;6.

## Figure Legends

38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51 Figure 1. Scattered Syndecan-1 positive cells are found in the stroma of prostate  
52 carcinoma tissue. (A) Expression of Syndecan-1 in adjacent normal and prostate  
53 adenocarcinoma tissue. Syndecan-1 is expressed in the basal cells of adjacent normal tissue.  
54  
55  
56  
57  
58 In prostate carcinoma Syncecan-1 expression is lost or found to be broadly expressed in the  
59  
60

1  
2  
3 epithelium. In addition, in a subset of prostate cancer cases Syndecan-1 positive cells were  
4  
5 found scattered in the stroma (shown in the insets and by arrows), but not in adjacent  
6  
7 normal prostate tissue. We call these cells Prostate Cancer Syndecan-1 Positive (PCSP) cells  
8  
9 and they could be found in the contexts of both epithelial Syndecan-1 alterations in tumors,  
10  
11 including expression across all epithelial cells, and where Syndecan-1 expression is lost in  
12  
13 the epithelium. (B) Examples of PCSP cells detected by anti-Syndecan-1 IHC. PCSP cells were  
14  
15 located in the stroma and tend to appear near regions that appear to contain immune cells  
16  
17 with small, round, hyperchromatic nuclei. Images are from 3 Independent patient cohorts.  
18  
19 Nuclei are counterstained with hematoxylin (blue) in IHC, or with DAPI (blue) in IF. IHC scale  
20  
21 bars – 100um; IF scale bars – 50um. SDC1 – Syndecan-1.

22  
23  
24  
25  
26  
27 Figure 2. Three distinct anti-Syndecan-1 antibodies show the expected patterns of  
28  
29 Syndecan-1 expression in control prostate and tonsil tissue, and stain scattered stromal  
30  
31 PCSP cells. (A) Staining of adjacent tissue sections of a representative adjacent normal  
32  
33 prostate and tonsil positive controls with two different mouse monoclonal antibodies  
34  
35 recognizing Syndecan-1. Both antibodies stained basal cells in adjacent normal prostate  
36  
37 epithelium and epithelium in tonsil as expected. (B) Staining of adjacent tissue sections of  
38  
39 representative adjacent normal prostate and tonsil positive controls with a mouse  
40  
41 monoclonal and rabbit polyclonal antibodies that both recognize Syndecan-1. These  
42  
43 antibodies also stained basal cells in adjacent normal prostate epithelium and epithelium in  
44  
45 tonsil as expected. (C) Three anti-Syndecan-1 antibodies recognise similar stromally-located  
46  
47 PCSP cells in both prostate tumors and in adjacent normal prostate tissue. Example cells are  
48  
49 highlighted in the magnified insets and with the arrows. Nuclei are counterstained with  
50  
51 hematoxylin (blue) for IHC. Scale bars – 100um with inserts at 3x zoom; SDC1 – Syndecan-1;  
52  
53 mo – mouse; rb – rabbit.  
54  
55  
56  
57  
58  
59  
60



1  
2  
3 Figure 3. Staining from two anti-Syndecan-1 antibodies overlap in individual  
4 prostate epithelial cells and in the PCSP cells seen in the stroma of prostate tumors. Mouse  
5 monoclonal anti-Syndecan-1 antibody #1 and the rabbit anti-Syndecan-1 antibodies stained  
6 the same cell population following IF staining of prostate tissue. Overlap was seen in the  
7 basal cells of adjacent normal (Upper panels: See inserts) and in the PCSP cells found in the  
8 stroma of tumor tissues (Lower panels: See arrows and inserts). Nuclei were counterstained  
9 with DAPI (blue) in IF. Scale bars – 50um. SDC1 – Syndecan-1; mo – mouse; rb – rabbit.

10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20 Figure 4. PCSP cells do not express common markers of immune cell lineages. FFPE  
21 sections of prostate tumor tissue were stained for Syndecan-1 (green) and immune cell  
22 markers (red). Nuclei were counterstained with DAPI (blue). (A) PCSP cells (inset) do not  
23 express the hematopoietic lineage marker CD45, the B-cell marker CD19, the plasma cell  
24 marker CD27 or the lymphocyte and mesenchymal stem cell marker CD73. (B)  
25  
26  
27  
28  
29  
30  
31  
32 Quantification of the percentage overlap of Syndecan-1+ cells with the marker stains.  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
Counts of PCSP cells were pooled from a minimum of 3 different patient samples, with a  
minimum of 200 cells counted in total per marker over at least 5 fields of view, with the  
exception of CD73 where 58 PCSP cells were counted. Scale bars – 50um. Insets: individual  
PCSP cells displayed at 3x zoom. SDC1 – Syndecan-1.

Figure 5. PCSP cells do not express markers of epithelial and secretory prostate cells.  
FFPE sections of prostate tumor tissue were stained for Syndecan-1 (green) and epithelial  
markers (red). Nuclei were counterstained with DAPI (blue). (A) PCSP cells (inset) do not  
express the epithelial cell markers pan-Cytokeratin or E-Cadherin. Similarly, the secretory  
epithelial marker PSA was not expressed in PCSP cells. (B) Quantification of the percentage  
overlap of PCSP cells with marker stains. Counts of PCSP cells were pooled from a minimum  
of 3 different patient samples. A minimum of 200 cells were counted in total per marker



1  
2  
3 over at least 5 fields of view, with the exception of E-Cadherin where 176 cells were  
4  
5 counted. Insets: individual PCSP cells displayed at 3x zoom. (C) Sequential sections of  
6  
7 adjacent normal prostate and prostate tumors were stained for Syndecan-1 and the basal  
8  
9 cell markers p63 and CK5, showing no colocalization in PCSP cells of tumor cases, despite  
10  
11 the overlap seen in the basal epithelial cells of adjacent normal tissue. Insets and arrows  
12  
13 show representative PCSP cells. Scale bars – 50um. SDC1 – Syndecan-1.  
14  
15  
16

17 Figure 6. PCSP cells do not express common markers consistent with a mesenchymal,  
18  
19 stromal, neural or endothelial cell identity. FFPE sections of prostate tumor tissue were  
20  
21 stained for Syndecan-1 (green) and mesenchymal/stromal cell type markers (red), and  
22  
23 nuclei were counterstained with DAPI (blue). (A) Mesenchymal cell type markers Vimentin,  
24  
25 N-Cadherin, nerve marker S100 and endothelial cell marker CD31. (B) Quantification of the  
26  
27 percentage overlap of PCSP cells with marker stains. Counts of PCSP cells were pooled from  
28  
29 a minimum of 3 different patient samples. A minimum of 200 cells were counted in total per  
30  
31 marker over at least 5 fields of view, with the exception of Vimentin, N-Cadherin and S100,  
32  
33 where 69, 197 and 29 cells were counted respectively. Insets: individual PCSP stromal cells  
34  
35 displayed at 3x zoom. Scale bars – 50um. SDC1 – Syndecan-1.  
36  
37  
38  
39  
40

41 Figure 7. PCSP cells are elongated and have prominent lamellipodia-like structures.  
42  
43 (A) Pooled single cell measurement data of PCSP cells from both tissue microarray cohorts.  
44  
45 A continuum between a more rounded cell morphology (circularity and aspect ratio close to  
46  
47 1) and a tail of cells assuming a more elongated cell shape with lower circularity and higher  
48  
49 aspect ratio was observed. (B) Sequential slices through a z-stack of images of PCSP cells in a  
50  
51 prostate tumor tissue section. PCSP cells had an elongated cell morphology, with polarized  
52  
53 nuclei and lamellipodia-like protrusions. Optical sections are 1um apart. Blue: DAPI; Green:  
54  
55 Syndecan-1. Scale bar: 10um. SDC1 – Syndecan-1.  
56  
57  
58  
59  
60

1  
2  
3 Figure 8. Distribution of PCSP cells in normal, adjacent normal and prostate  
4 carcinoma. (A) Plots of PCSP cell burden in cohort 1. PCSP cells were detected in 0/8 normal  
5  
6 carcinoma. (A) Plots of PCSP cell burden in cohort 1. PCSP cells were detected in 0/8 normal  
7  
8 samples, 4/8 adjacent normal (NAT) samples and 28/80 malignant samples. (B) Plots of PCSP  
9  
10 cell burden in cohort 1 and (C) cohort 2 malignant tissues showed that a subset of Gleason 5  
11  
12 patients had a particularly high burden in both cohorts.  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60




Figure 1. Scattered Syndecan-1 positive cells are found in the stroma of prostate carcinoma tissue. (A) Expression of Syndecan-1 in adjacent normal and prostate adenocarcinoma tissue. Syndecan-1 is expressed in the basal cells of adjacent normal tissue. In prostate carcinoma Syndecan-1 expression is lost or found to be broadly expressed in the epithelium. In addition, in a subset of prostate cancer cases Syndecan-1 positive cells were found scattered in the stroma (shown in the insets and by arrows), but not in adjacent normal prostate tissue. We call these cells Prostate Cancer Syndecan-1 Positive (PCSP) cells and they could be found in the contexts of both epithelial Syndecan-1 alterations in tumors, including expression across all epithelial cells, and where Syndecan-1 expression is lost in the epithelium. (B) Examples of PCSP cells detected by anti-Syndecan-1 IHC. PCSP cells were located in the stroma and tend to appear near regions that appear to contain immune cells with small, round, hyperchromatic nuclei. Images are from 3 independent patient cohorts. Nuclei are counterstained with hematoxylin (blue) in IHC, or with DAPI (blue) in IF. IHC scale bars – 100um; IF scale bars – 50um. SDC1 – Syndecan-1.

278x293mm (300 x 300 DPI)




Figure 2. Three distinct anti-Syndecan-1 antibodies show the expected patterns of Syndecan-1 expression in control prostate and tonsil tissue, and stain scattered stromal PCSP cells. (A) Staining of adjacent tissue sections of a representative adjacent normal prostate and tonsil positive controls with two different mouse monoclonal antibodies recognizing Syndecan-1. Both antibodies stained basal cells in adjacent normal prostate epithelium and epithelium in tonsil as expected. (B) Staining of adjacent tissue sections of representative adjacent normal prostate and tonsil positive controls with a mouse monoclonal and rabbit polyclonal antibodies that both recognize Syndecan-1. These antibodies also stained basal cells in adjacent normal prostate epithelium and epithelium in tonsil as expected. (C) Three anti-Syndecan-1 antibodies recognise similar stromally-located PCSP cells in both prostate tumors and in adjacent normal prostate tissue. Example cells are highlighted in the magnified insets and with the arrows. Nuclei are counterstained with hematoxylin (blue) for IHC. Scale bars – 100um with inserts at 3x zoom; SDC1 – Syndecan-1; mo – mouse; rb – rabbit.

494x332mm (300 x 300 DPI)





Figure 3. Staining from two anti-Syndecan-1 antibodies overlap in individual prostate epithelial cells and in the PCSP cells seen in the stroma of prostate tumors. Mouse monoclonal anti-Syndecan-1 antibody #1 and the rabbit anti-Syndecan-1 antibodies stained the same cell population following IF staining of prostate tissue. Overlap was seen in the basal cells of adjacent normal (Upper panels: See inserts) and in the PCSP cells found in the stroma of tumor tissues (Lower panels: See arrows and inserts). Nuclei were counterstained with DAPI (blue) in IF. Scale bars – 50um. SDC1 – Syndecan-1; mo – mouse; rb – rabbit.

33 352x205mm (300 x 300 DPI)





33 Figure 4. PCSP cells do not express common markers of immune cell lineages. FFPE sections of prostate  
 34 tumor tissue were stained for Syndecan-1 (green) and immune cell markers (red). Nuclei were  
 35 counterstained with DAPI (blue). (A) PCSP cells (inset) do not express the hematopoietic lineage marker  
 36 CD45, the B-cell marker CD19, the plasma cell marker CD27 or the lymphocyte and mesenchymal stem cell  
 37 marker CD73. (B) Quantification of the percentage overlap of Syndecan-1+ cells with the marker stains.  
 38 Counts of PCSP cells were pooled from a minimum of 3 different patient samples, with a minimum of 200  
 39 cells counted in total per marker over at least 5 fields of view, with the exception of CD73 where 58 PCSP  
 40 cells were counted. Scale bars – 50um. Insets: individual PCSP cells displayed at 3x zoom. SDC1 –  
 41 Syndecan-1.

42 295x226mm (300 x 300 DPI)

43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60




Figure 5. PCSP cells do not express markers of epithelial and secretory prostate cells. FFPE sections of prostate tumor tissue were stained for Syndecan-1 (green) and epithelial markers (red). Nuclei were counterstained with DAPI (blue). (A) PCSP cells (inset) do not express the epithelial cell markers pan-Cytokeratin or E-Cadherin. Similarly, the secretory epithelial marker PSA was not expressed in PCSP cells. (B) Quantification of the percentage overlap of PCSP cells with marker stains. Counts of PCSP cells were pooled from a minimum of 3 different patient samples. A minimum of 200 cells were counted in total per marker over at least 5 fields of view, with the exception of E-Cadherin where 176 cells were counted. Insets: individual PCSP cells displayed at 3x zoom. (C) Sequential sections of adjacent normal prostate and prostate tumors were stained for Syndecan-1 and the basal cell markers p63 and CK5, showing no colocalization in PCSP cells of tumor cases, despite the overlap seen in the basal epithelial cells of adjacent normal tissue. Insets and arrows show representative PCSP cells. Scale bars – 50um. SDC1 – Syndecan-1.

325x335mm (300 x 300 DPI)




Figure 6. PCSP cells do not express common markers consistent with a mesenchymal, stromal, neural or endothelial cell identity. FFPE sections of prostate tumor tissue were stained for Syndecan-1 (green) and mesenchymal/stromal cell type markers (red), and nuclei were counterstained with DAPI (blue). (A) Mesenchymal cell type markers Vimentin, N-Cadherin, nerve marker S100 and endothelial cell marker CD31. (B) Quantification of the percentage overlap of PCSP cells with marker stains. Counts of PCSP cells were pooled from a minimum of 3 different patient samples. A minimum of 200 cells were counted in total per marker over at least 5 fields of view, with the exception of Vimentin, N-Cadherin and S100, where 69, 197 and 29 cells were counted respectively. Insets: individual PCSP stromal cells displayed at 3x zoom. Scale bars – 50µm. SDC1 – Syndecan-1.

277x192mm (300 x 300 DPI)







Figure 7. PCSP cells are elongated and have prominent lamellipodia-like structures. (A) Pooled single cell measurement data of PCSP cells from both tissue microarray cohorts. A continuum between a more rounded cell morphology (circularity and aspect ratio close to 1) and a tail of cells assuming a more elongated cell shape with lower circularity and higher aspect ratio was observed. (B) Sequential slices through a z-stack of images of PCSP cells in a prostate tumor tissue section. PCSP cells had an elongated cell morphology, with polarized nuclei and lamellipodia-like protrusions. Optical sections are 1 $\mu$ m apart. Blue: DAPI; Green: Syndecan-1. Scale bar: 10 $\mu$ m. SDC1 – Syndecan-1.

196x231mm (300 x 300 DPI)





Figure 8. Distribution of PCSP cells in normal, adjacent normal and prostate carcinoma. (A) Plots of PCSP cell burden in cohort 1. PCSP cells were detected in 0/8 normal samples, 4/8 adjacent normal (NAT) samples and 28/80 malignant samples. (B) Plots of PCSP cell burden in cohort 1 and (C) cohort 2 malignant tissues showed that a subset of Gleason 5 patients had a particularly high burden in both cohorts.

263x128mm (300 x 300 DPI)



**Supplementary Figure 1.** Immunohistochemistry with anti-immune cell marker antibodies on positive control paraffin-embedded tissue sections. As expected, CD45, CD19, CD27 showed cytoplasmic and membranous staining in lymphocyte populations in the tonsil and CD73 marked lymphocytes in the placenta. Nuclei were counterstained with haematoxylin (blue). IHC scale bars – 100um.

	Cohort 1 (PR1921)		Cohort 2 (PR803b)	
		%		%
Number of Patients	80		71	
<b>Age:</b>				
Range	20-85		20-87	
Mean	67.85		67.6	
Median	69.5		70	
<50	2	2.5	3	4.2
50-59	5	6.3	4	5.6
60-69	33	41.3	27	38.0
>=70	40	50.0	37	52.1
<b>Clinical Stage:</b>				
I	6	7.5	2	2.8
II	37	46.3	33	46.5
III	14	17.5	10	14.1
IV	22	27.5	26	36.6
ND	1	1.3	0	0.0
<b>Primary Gleason Grade:</b>				
3	8	10.0	19	26.8
4	36	45.0	24	33.8
5	30	37.5	24	33.8
ND	6	7.5	4	5.6
<b>Overall Gleason Grade:</b>				
<=Gleason 3+4	8	10.0	16	22.5
>=Gleason 4+3	66	82.5	51	71.8
ND	6	7.5	4	5.6
<b>T Category:</b>				
T1	4	5.0	2	2.8
T2	47	58.8	40	56.3
T3	22	27.5	21	29.6
T4	6	7.5	8	11.3
ND	1	1.3	0	0.0
<b>N Category:</b>				
N0	64	80.0	51	71.8
N1	15	18.8	18	25.4
N2		0.0	2	2.8
ND	1	1.3	0	0.0

M Category				
M0	64	80.0	49	69.0
M1	15	18.8	22	31.0
ND	1	1.3	0	0.0

SDC1 Epithelial Reactivity:

Absent	66	82.5	57	80.3
Present	14	17.5	13	18.3
ND	0	0.0	1	1.4

PCSP Cell Presence:

Absent	52	65.0	56	78.9
Present	28	35.0	15	21.1

Pre Peer Review

Antibody	Species of Origin	Clonality [Clone Number]	Working Dilution	Supplier	Catalogue No.
Syndecan-1 (SDC1)	Mouse	Monoclonal [B-A38]	1:200	Novus Biologicals	NB100-64980
Syndecan-1 (SDC1)	Rabbit	Polyclonal	1:50	Santa-Cruz Biotechnology	Sc-5632
Syndecan-1 (SDC1)	Mouse	Monoclonal [MI15]	1:25	Dako	N/A
CD45	Rabbit	Polyclonal	1:50	Santa-Cruz Biotechnology	Sc-25590
CD19	Rabbit	Monoclonal [EPR5906]	1:250	Abcam	Ab134114
CD27	Rabbit	Monoclonal [EPR8569]	1:100	Abcam	Ab131254
CD73	Rabbit	Polyclonal	1:25	Abcam	Ab115289
Pan-Cytokeratin	Mouse	Monoclonal [C-11]	1:100	Sigma-Aldrich	C2931
E-Cadherin	Mouse	Monoclonal [4A2C7]	1:25	Thermo Fisher Scientific	33-4000
Prostate-Specific Antigen (PSA)	Rabbit	Polyclonal	1:50	Dako	A0562
p63	Mouse	Monoclonal [BC4A4]	1:50	Abcam	Ab735
Cytokeratin -5	Rabbit	Polyclonal	1:1000	Abcam	Ab24647
Vimentin	Mouse	Monoclonal [V9]	1:50	Santa-Cruz Biotechnology	Sc-620

---

N-Cadherin	Rabbit	Polyclonal	1:100	Abcam	Ab18203
S100	Rabbit	Polyclonal	1:2000	Dako	Z0311
CD31	Mouse	Monoclonal [JC70A]	1:30	Dako	M0823

---

For Peer Review

	Cohort 1 (PR1921)		Cohort 2 (PR803b)	
	Mean across samples	Stdev across samples	Mean across samples	Stdev across samples
<b>Area</b>	66.42	35.09	72.14	41.31
<b>Circularity</b>	0.69	0.12	0.70	0.13
<b>Aspect Ratio</b>	1.88	0.62	1.83	0.68
<b>Roundness</b>	0.58	0.16	0.60	0.17
<b>Solidity</b>	0.88	0.06	0.88	0.07