



Republic Of Iraq
Ministry of higher Education and
Scientific research University Of
Thi-Qar College Of Medicine



Association Between Nausea and Vomiting During Pregnancy and Sleep Quality

A Project Submitted to
the College of Medicine, University of Thi-Qar,
in Partial Fulfillment of the Requirement For The Degree & Bachelor of Science in
Medicine.

Supervised by :

Dr. Alaa Hussein Ali Al-naser

Dne by:

Nabaa Ahmed Sattar

Hawraa Haider jabbar

Manal Lateef Hussain

Rawan Haider Ibrahim

DEDICATION

Thankfully to Almighty God, for guidance, power , and protection and for giving us a healthy life to be able to carry out this research.

This study is wholeheartedly dedicated to our beloved parents, who have been our source of inspiration and strength throughout this research. Their continuous moral, spiritual, emotional, and financial support to us has helped us finish this research. All of these, we offer to you.

And lastly, we dedicate this research to our adviser, For her tremendous effort and patience to guide us throughout the research and helped us to better understand the key points of this research. Furthermore, to our classmates who shared their words of advice and encouragement that helped us be motivated to finish this research.

Purpose: Nausea and vomiting during pregnancy (NVP) of varying degrees of severity are commonly experienced by pregnant women. This paper explores the association between NVP and poor sleep quality.

Patients and Methods: A cross-sectional study was conducted in Bint Al Huda Hospital and an obstetrics clinic. A total of 125 pregnant women (representing a response rate of 88%) completed a self-administered questionnaire.

Results: Of the 125 participants, the mean sleep duration was 7.76 hours, and 55% of them report poor sleep quality. In this study, 50.2%, 48.1% and 1.7% women reported mild, moderate, and severe NVP, respectively. Compared with women with mild NVP, women with moderate or severe NVP were more likely to report poor sleep quality. After adjusted for demographics and gestational age, moderate and severe NVP were associated with poor sleep quality.

Conclusion: Our study suggested that moderate and severe NVP increase the risk of poor sleep quality. Further studies are warranted that focus on the mechanisms of the association between NVP and poor sleep quality.

Keywords: nausea, vomiting, pregnancy, sleep quality.

Introduction

Nausea and vomiting are most commonly experienced during early pregnancy. In different samples from prior studies, the prevalence of nausea and vomiting of pregnancy (NVP) has been found to range from 35% to 91% . [1-3] However, NVP symptoms are often ignored by pregnant women and clinicians, or pregnant women may refuse treatment for NVP because of concerns about the adverse effects of any treatment on their pregnancy. However, if NVP deteriorates into its severest form, hyperemesis gravidarum, it can threaten both mother and fetus. [6–8]

Sleep quality typically decreases during pregnancy. Furthermore, it has been found that sleep quality are more likely to undergo preterm labor[10] or a caesarean delivery.[11] Previous research has also shown that poor sleep quality during pregnancy increases the risk of perinatal depression[12] and suicide.[13] NVP may be an important risk factor for reduction in sleep quality. [14] while a case-control study conducted among pregnant women reported that hyperemesis gravidarum likewise increased the risk of poor sleep quality.[15] However, most previous research has been carried out with reference to patients with hyperemesis gravidarum, and only a few studies to date have investigated the association between NVP and sleep quality in pregnant women who do not completely meet the diagnostic criteria of hyperemesis gravidarum.

Stressors and mental health status play important roles with regard to poor sleep quality during pregnancy. Specifically, somatic and psychological stressors during pregnancy can activate stress systems, such as a hypothalamic–pituitary–adrenal (HPA) axis stress response and inflammatory reaction.[16] Prior research has concluded that NVP, as a somatic stressor, increases the risk of mental health problems such as depression and anxiety.[8,17] Equally, depressive symptoms have been found to increase the risk of poor sleep quality during pregnancy. We hypothesize that severe NVP increases the risk of poor sleep quality for pregnant women. To assess this postulation, we conducted a cross-sectional study to examine the association between NVP and poor sleep quality

Patients and Methods:

Participants

All participants were recruited from the Bint Al Huda Hospital and Gyne and Obstetric clinic of DhiQar .Pregnant women were eligible to participate if they were 16 years of age or older, were between weeks 6 and 28 of their pregnancy, and they planned to attend perinatal care and deliver at this hospital. With the help of trained interviewers, each woman completed a self-administered questionnaire. A total of 125 women completed the questionnaire (representing a response rate of 88%). All data were collected between March 2022and May 2022. All participants provided written informed consent.

Measures:

NVP

NVP was measured using the modified Pregnancy-Unique Quantification of Emesis and Nausea (PUQE) index.[19] Compared to the original PUQE index, which focused on symptoms experienced during the preceding the previous 24 hours [20] the modified PUQE covers the first trimester of pregnancy. It presents three subscales (questions referring to nausea, vomiting, and retching, respectively), and the score for each subscale ranges from 1 to 5 points. Thus, the total score from the modified PUQE index ranges from 3 to 15 points, with a higher score indicating more severe NVP. Lacasse et al provided the recommended cut-off total scores corresponding to “mild NVP” (≤ 6 points), “moderate NVP” (7–12 points), and “severe NVP” (≥ 13 points) in relation to the modified PUQE index.

Sleep Quality

The Pittsburgh Sleep Quality Index (PSQI) was used to measure participants' sleep quality over the preceding 30 days.[22] The duration of each night's sleep was also calculated women who experience poor

Across the PSQI's seven components, responses to each of which are rated from 1 (“no difficulty”) to 3 (“severe difficulty”), the overall score can range from 0 to 21 points, with higher scores indicating poorer sleep .In pregnant women, the cut-off score for identifying cases of poor sleep quality was 5.

Covariates

Demographical and gestational age data that have been reported as being associated with NVP or poor sleep quality were considered as covariates in the present study. Demographical data encompassing maternal age, education level, work status, household income per capita, and marital status were recorded via self-report. For example, education level was measured by asking each participant to specify the highest level of education that they had attained (“junior high school or below,” “senior high school,” or “college or university”). Work status was measured by asking participants if they were working, including full-time and part-time jobs, with their responses being either “working” or “not working.” Household income per capita was calculated by dividing the total household income in the last year by the household’s population. Marital status was measured by asking about the participant’s current marital status, with their responses categorized as “married” or “unmarried or divorced.”

Statistical Analysis

Sleep quality, NVP, depressive symptoms, and demographic data were assessed using frequency tables for categorical variables and compared using Pearson’s chi-squared test. Maternal age and gestational age were summarized using means and standard deviations (SDs), and compared using a student’s t-test. According to the recommended cut-off scores of the modified PUQE index, participants were divided into three groups (corresponding to mild NVP, moderate NVP, and severe NVP), while their PSQI scores

were summarized with means and SDs and compared using analysis of variance. Logistic regression analysis was used to assess the association between NVP and sleep quality, and odds ratios (ORs) were obtained with 95% confidence intervals (CIs). The variables that were widely reported in previous studies or that were significant at the 0.10 level in the univariate logistic regression were entered into the multivariate logistic regression models. Three sets of logistic regression models were performed to evaluate the hypothesized associations. First, a univariate logistic regression model (Model 1) was run without being adjusted for the covariates. In Model 2, demographic data and gestational age were adjusted. Next, in Model 3, depressive symptoms and the covariates of Model 2 were adjusted. Structural equation modelling was used to measure the mediating effect of depressive symptoms on the association between NVP and poor sleep quality. The structural equation modelling was performed using the robust maximum-likelihood estimation. Standardized coefficients and indirect and total effects were estimated, and the 95% CI was measured using the bootstrap method with 2000 resamplings.

Results

Participants

Participants' demographic characteristics are detailed in Table 1. Of the 2494 participants who were analyzed, the mean age was 29.6 years old (SD = 4.8, range 18–46), and the mean gestational age was 22.9 (SD = 10.1, range 9–36) weeks. About half of the participants had graduated from college or university (52.0%). Most of the participants were working (75.0%) and were married (95.4%). NVP

The mean PUQE score of participants was 6.65 (SD = 2.56, range 3–15). According to the definitions of the modified PUQE, in this study, 49.1% 49.3% and 1.6% women were evaluated to be experiencing mild, moderate and severe NVP, respectively.

Sleep Quality

The mean sleep duration of participants was 7.76 (SD = 1.38, range 3.00–13.00) hours, and 83.2% participants sleep more 7 hours each day. The means global PSQI score was 6.39 (SD = 3.18, range 0–17), and 54.3% reported poor sleep quality (PSQI score > 5). First trimester (53.8%) and third trimester (60.8%) pregnant women were more likely to report poor sleep quality than second trimester pregnant women (48.3% $\chi^2 = 29.39$, $p < 0.001$). Compared with working women, non-working women were more likely to report poor sleep quality ($\chi^2 = 8.69$, $p = 0.003$)

Table 1 Characteristics of Participants and Sleep Quality (N= 125)

Characteristics	Total n (%)	Poor Sleep Quality ^a	
		With n (%)	Without n (%)
Total	125 (100)	67 (54.3)	57 . (45.7)
Gestational age			
First trimester	31 (24.8)	67 (53.8)	57 (46.2)
Second trimester	47 (37.8)	60 (48.3)	65 (51.7)
Third trimester	46 (37.4)	76 (60.8)	49 (39.2)
Education level			
Junior high school or below	30 (24.3)	67 (53.4)	58 (46.6)
Senior high school	29 (23.4)	60 (48.3)	61 (48.6)
College or university	65 (52)	70 (56.2)	55 (43.8)
Missing data	1 (0.8)	NA	NA
Work status			
Not working	30 (24.4)	362 (59.4)	247 (40.6)
Working	94 (75)	984 (52.6)	362 (47.4)

Depressive symptoms

The mean EPDS score of participants was 7.39 (SD = 4.14, range 0–28). There were 34.6% participants who reported depressive symptoms (EPDS score ≥ 9). Women who were enduring poor sleep quality (67.9%) were more likely to report depressive symptoms than women who had not experienced poor sleep quality (32.1% $\chi^2 = 93.43$, $p < 0.001$). Women with moderate (39.8%) and severe NVP (40.0%) were more likely to report depressive symptoms than women with mild NVP (29.2% $\chi^2 = 31.08$, $p < 0.001$).

Association Between NVP and Sleep quality

Table 2 presents the relative sleep quality of participants alongside their varying degrees of NVP. As shown, there was no significant difference in sleep duration among women with different degrees of NVP ($F = 0.61$, $p = 0.545$). However, the global PSQI scores were higher in women with severe NVP than for women with mild or moderate NVP ($F = 22.04$, $p < 0.001$). Of the seven PSQI components, women with severe NVP were higher in their scores for “subjective sleep quality”, “sleep latency”

N=125 , PSQ =>5

Number	%	Sleep (h/day)	Quality
104	83.2	>7	Good
68	54.3	>5	Poor
First trimester 67	53.8	4-5	Poor
Second trimester 60	48.3	>5	Good
Third trimester 76	60.8	4-5	Poor

habitual sleep efficiency”, “sleep disturbance”, and “day- time dysfunction” than were women with mild or moderate NVP ($p < 0.05$). There was no significant difference regarding the PSQI components of “sleep duration” and “use of sleep medication” among women with different degrees of NVP ($p > 0.05$). The prevalence of poor sleep quality among women who reported mild, moderate, and severe NVP was 49.0%, 59.0% and 72.5% respectively. Compared with women with mild NVP, women with moderate or severe NVP were more likely to report poor sleep quality ($\chi^2 = 30.16, p < 0.001$).

Logistic regression models were conducted to assess the association between NVP and sleep quality (see Table 3). Without adjusting any covariate (Model 1), compared with women with mild NVP, women with moderate or severe NVP were at higher risk of poor sleep

quality: crude OR (COR) = 1.50, 95%CI = 1.28–1.76 and COR = 2.74, 95% CI = 1.36–5.53, respectively. After adjusting for maternal age, gestational age, educational level, work status, household income per capita, and marital status (Model 2), moderate and severe NVP were still associated with poor sleep quality: adjusted OR (AOR) = 1.66, 95% CI = 1.40–1.96, and AOR = 2.95, 95%CI = 1.44–6.02, respectively. In Model 2, we also examined the moderating effect of gestational age on the association between NVP and sleep quality. However, as there was found to be no significant moderating effect of gestational age ($p = 0.125$), we inferred that there was no difference in the association between NVP and poor sleep quality across the three trimesters. In Model 3, after further adjusting for depressive symptoms, NVP was still found to be associated with poor sleep quality, but the extent of this association was weaker.

Discussion

Nausea and vomiting bring a substantial health burden to pregnant women, and can reduce their overall quality of life. Poor sleep quality is also a significant health problem during pregnancy that can increase the risk of pregnancy complication, including gestational diabetes mellitus²⁶ and gestational hypertension.²⁷ In this study, we found that severe NVP was associated with poor sleep quality. In our study, 54.3% of the sample of pregnant women self-reported poor sleep quality. This result is consistent with the findings of previous research,^{9,26} which collectively suggests that poor sleep quality is an issue for Chinese pregnant women. We found that there was no difference in sleep

duration between pregnant women who reported mild, moderate, or severe NVP.

However, NVP was found to be associated with poor sleep quality. Specifically, compared with women who reported mild NVP, those who reported moderate or severe NVP were more likely to suffer with poor sleep quality. This finding, too, is similar to those of previous studies. For example, Swallow et al reported that severe NVP increases the risk of insomnia among early pregnant women.²⁸ Yildırım et al conducted a casecontrol study among 98 pregnant women, and reported that, compared with a health control group (7.7%), women who were diagnosed with hyperemesis gravidarum were more likely to report poor sleep quality (60.9%).¹⁵ In fact, our findings suggest that poor sleep quality is not limited to women with severe NVP who meet the diagnostic criteria of hyperemesis gravidarum, but is also found in women with moderate NVP. The biological mechanism of NVP is still not fully understood. Previous studies have suggested that the condition is associated with increased levels of reproductive hormones during pregnancy, such as estrogen and progesterone²⁹ and these hormones have also been found to be linked to sleep quality.³⁰ Prior research has shown that high levels of estradiol increase the risk of restless legs syndrome and poor sleep quality among pregnant women.³¹ Also, in other animal studies, estrogen and progesterone have been observed to decrease the duration of non-rapid eye movement sleep and rapid eye movement sleep.³² Such evidence may partially explain the mechanism of the association between NVP and poor sleep quality, but more research is needed. Previous studies have

also shown that poor sleep quality is associated with gestational age and that poor sleep quality is more likely to be reported in the first and third trimesters.^{33,34} Our results in this regard are broadly consistent

with those of prior research. However, in the present study, the moderating effect of gestational age on the association between NVP and sleep quality was not found to be significant. Such a finding suggested that there might be a long-term effect of NVP on sleep quality. Therefore, we further explored the role of mental health status on the association between NVP and poor sleep quality. In addition, poor sleep quality is associated with mental health problems among pregnant women. A prospective longitudinal study reported that anxiety and depression is associated with poor sleep quality, and that this association is significant throughout the pregnancy.¹² Most depressed patients experience disturbances of the HPA axis, such as hypercortisolemia and glucocorticoid receptor expression reduction.⁴⁰ Numerous studies have suggested that disturbance of the HPA axis may be an important mechanism of sleep disorders in both pregnant³³ and non-pregnant individuals.⁴¹ The HPA axis's functions change during pregnancy, including a rise in cortisol concentrations rise and a blunting of diurnal rhythm.⁴² This in turn means that depressive symptoms during pregnancy may be more likely to cause sleep problems than in non-pregnant women. Therefore, depressive symptoms may at least partially explain the long-term effect of NVP on sleep quality. Our findings suggest that NVP is associated with poor sleep quality, and this result has clinical implications because both NVP and sleep quality are modifiable factors. On the one hand, in perinatal care, more attention

should be paid to screening for sleep problems and depression, especially in women with severe NVP. On the other hand, for women with severe NVP, effective interventions such as ginger, vitamin supplementation, fluid rehydration, and drug

therapy should be taken to reduce NVP symptoms,²⁹ and thereby potentially improve sleep quality during pregnancy. The present study features several strengths, including, first, the fact that it was conducted among pregnant women, and not limited to hyperemesis gravidarum patients, which allowed us to explore the association between mild or moderate NVP and poor sleep quality. Second, sleep quality and NVP were measured using the PSQI and the modified PUQE, respectively, which have been verified in terms of their reliability and validity. There are also several limitations to our study that should be noted as well. First, given the research's cross-sectional study design, it is difficult to make causal inferences. Second, as sleep quality was measured using a self-reporting questionnaire, rather than through objective records, recall bias cannot be completely ruled out. Third, due to the limitation of the study's out-patient clinic setting, data pertaining to potential confounding factors such as physical exercise and smoking were not collected and corrected in the multivariate analysis. Conclusion Our study's findings suggest that moderate and severe NVP increases the risk of poor sleep quality among Chinese pregnant women. Moreover, depressive symptoms were found to mediate the association between NVP and poor sleep quality. Therefore, more attention should be given to NVP, depressive symptoms, and sleep quality in prenatal care, and screening and

intervention in respect of all three factors should be considered an important part of such care. Future research is needed in order to understand the mechanisms of the association between NVP and poor sleep quality. In particular, the psychological mediation processes associated with

NVP, such as depression, anxiety, and eating disorders, should be considered in future research.

Acknowledgments References

1. Chan OK, Sahota DS, Leung TY, Chan LW, Fung TY, Lau TK. Nausea and vomiting in healthrelated quality of life among Chinese pregnant women. *Aust NZ J Obstet Gynaecol.* 2010;50(6):512–518.

doi:10.1111/j.1479-828X.2010.01216.x submit your manuscript |

www.dovepress.com 47 DovePress 2. Einarson TR, Piwko C, Koren G.

Quantifying the global rates of nausea and vomiting of pregnancy: a meta analysis. *J Popul Ther Clin Pharmacol.* 2013;20(2):e171. 3. Fejzo

MS, Trovik J, Grooten IJ, et al. Nausea and vomiting of pregnancy and hyperemesis gravidarum. *Nat Rev Dis Primers.* 2019;5(1):62.

doi:10.1038/s41572-019-0110-3 4. Fossum S, Vikanes AV, Naess O, Vos

L, Grotmol T, Halvorsen S. Hyperemesis gravidarum and long-term mortality: a population-based cohort study. *BJOG.* 2017;124(7):1080–

1087. doi:10.1111/1471- 0528.14454 5. Mayer KH, McGill AL. Second-trimester fetal loss in a patient with hyperemesis gravidarum

complicated by refeeding syndrome. *Obstet Gynecol.*

2019;133(6):1167–1170. doi:10.1097/ AOG.000000 0000003276 6. Bai

G, Korfage IJ, Groen EH, Jaddoe VWV, Mautner E, Raat H. Associations between nausea, vomiting, fatigue and health-related quality of life of

women in early pregnancy: the Generation R study. *PLoS One*. 2016;11(11):e166133. doi:10.1371/journal.pone.0166133 7. Beyazit F, Sahin B. Effect of nausea and vomiting on anxiety and depression levels in early pregnancy. *Eurasian J Med*. 2018;50 (2):111–115. doi:10.5152/eurasianjmed.2018.170320 8. Bray N,

Grasby KL, Lind PA, Painter JN, Colodro-Conde L, Medland SE. The psychosocial impact of nausea and vomiting during pregnancy as a predictor of postpartum depression. *J Health Psychol*. 2019;1573745512. doi:10.1177/1359105319859048 9. Sedov ID,

Cameron EE, Madigan S, Tomfohr-Madsen LM. Sleep quality during pregnancy: a meta-analysis. *Sleep Med Rev*. 2018;38:168–176.

doi:10.1016/j.smrv.2017.06.005 10. Felder JN, Baer RJ, Rand L, Jelliffe-Pawlowski LL, Prather AA. Sleep disorder diagnosis during pregnancy and risk of preterm birth. *Obstet Gynecol*. 2017;130(3):573–581.

doi:10.1097/AOG.0000000000002132 11. Teong A, Diong AX, Omar SZ, Tan PC. The impact of self-reported sleep on caesarean delivery in women undergoing induction of labour: a prospective study. *Sci Rep*.

2017;7(1):12339. doi:10.1038/s41598-017-12410-7 12. Yu Y, Li M, Pu L, et al. Sleep was associated with depression and anxiety status during pregnancy: a prospective longitudinal study. *Arch Womens Ment Health*. 2017;20(5):695–701. doi:10.1007/s00737-017-0754-5 13.

Palagini L, Cipollone G, Masci I, et al. Stress-related sleep reactivity is associated with insomnia, psychopathology and suicidality in pregnant women: preliminary results. *Sleep Med*. 2019;56:145–150.

doi:10.1016/j.sleep.2019.01.009 14. Akyuz RG, Ugur O, Elcigil A. Sleep quality in lung cancer patients. *Asian Pac J Cancer Prev*.

2013;14(5):2909–2913. doi:10.7314/APJCP.2013.14.5.2909 15. Yildirim E, Demir E. The relationship of hyperemesis gravidarum with sleep disorders, anxiety and depression. *J Obstet Gynaecol*. 2019;39(6):793–798. doi:10.1080/01443615.2019.1572725 16. Palagini L, Gemignani A, Banti S, Manconi M, Mauri M, Riemann D. Chronic sleep loss during pregnancy as a determinant of stress:

impact on pregnancy outcome. *Sleep Med*. 2014;15 (8):853–859. doi:10.1016/j.sleep.2014.02.013 17. Kjeldgaard HK, Eberhard-Gran M, Benth J, Vikanes ÅV. Hyperemesis gravidarum and the risk of emotional distress during and after pregnancy. *Arch Womens Ment Health*. 2017;20 (6):747– 756. doi:10.1007/s00737-017-0770-5 18. Polo-Kantola P, Aukia L, Karlsson H, Karlsson L, Paavonen EJ. Sleep quality during pregnancy: associations with depressive and anxiety symptoms. *Acta Obstet Gynecol Scand*. 2017;96 (2):198–206. doi:10.1111/aogs.13056 19. Lacasse A, Rey E, Ferreira E, Morin C, Bérard A. Validity of a modified Pregnancy-Unique Quantification of Emesis and Nausea (PUQE) scoring index to assess severity of nausea and vomiting of pregnancy. *Am J Obstet Gynecol*. 2008;198(1):71. doi:10.1016/j.ajog.2007.05.051 20. Koren G, Magee L, Attard C, et al. A novel method for the evaluation of the severity of nausea and vomiting of pregnancy. *Eur J Obstet Gynecol Reprod Biol*. 2001;94(1):31–36. doi:10.1016/S0301-2115(00)00344-4 21. Yu-hua W, Lan W. Effect of health-related quality of life in women with nausea and vomiting of pregnancy (in Chinese). *Clin J Med Officers*. 2012;40(2):411–415. 22. Buysse DJ, Reynolds CFR, Monk TH, Berman SR, Kupfer DJ. The Pittsburgh Sleep Quality Index: a new instrument for psychiatric

practice and research. *Psychiatr Res.* 1989;28(2):193– 213.
doi:10.1016/0165-1781(89)90047-4 23. Cox JL, Holden JM, Sagovsky R.
Detection of postnatal depression. Development of the 10-item
Edinburgh Postnatal Depression Scale. *Br J Psychiatr.* 1987;150(6):782–
786. doi:10.1192/bjp.150.6.782 24. Howard LM, Ryan EG, Trevillion K,
et al. Accuracy of the Whooley questions and the Edinburgh Postnatal
Depression Scale in identifying depression and other mental

disorders in early pregnancy. *Br J Psychiatr.* 2018;212(1):50–56.
doi:10.1192/bjp.2017.9 25. Zhao Y, Kane I, Wang J, Shen B, Luo J, Shi S.
Combined use of the postpartum depression screening scale (PDSS) and
Edinburgh post- natal depression scale (EPDS)

to identify antenatal depression among Chinese pregnant women with
obstetric complications. *Psychiatr Res.* 2015;226(1):113–119.
doi:10.1016/j. psychres.2014.12.016 26. Wang H, Leng J, Li W, et al.
Sleep duration and quality, and risk of gestational diabetes mellitus in
pregnant Chinese women. *Diabetic Med.* 2017;34(1):44–50.
doi:10.1111/dme.13155 27. Dunietz GL, Shedden K, Lisabeth LD,
Treadwell MC, O'Brien LM. Maternal weight, snoring, and
hypertension: potential pathways of associations. *Am J Hypertens.*
2018;31(10):1133–1138. doi:10.1093/ajh/ hpy085 28. Swallow BL,
Lindow SW, Masson EA, Hay DM. Psychological health in early
pregnancy: relationship with nausea and vomiting. *J Obstet Gynaecol.*
2009;24(1):28–32. doi:10.1080/01443610310001620251 29. Bustos M,
Venkataramanan R, Caritis S. Nausea and vomiting of pregnancy -
What's new? *Auton Neurosci.* 2017;202:62–72.
doi:10.1016/j.autneu.2016.05.002 30. Carrier J, Semba K, Deurveilher S,

et al. Sex differences in age-related changes in the sleepwake cycle. *Front Neuroendocrin.* 2017;47:66–85. doi:10.1016/j.yfrne.2017.07.004

31. Dzaja A, Wehrle R, Lancel M, Pollmacher T. Elevated estradiol plasma levels in women with restless legs during pregnancy. *Sleep.* 2009;32(2):169–174. doi:10.1093/sleep/32.2.169

32. Deurveilher S, Rusak B, Semba K. Female reproductive hormones alter sleep architecture in ovariectomized rats. *Sleep.* 2011;34 (4):519–530.

doi:10.1093/sleep/34.4.519

33. Bublitz MH, Bourjeily G, D'Angelo C, Stroud LR. Maternal sleep quality and diurnal cortisol regulation over pregnancy. *Behav Sleep Med.* 2018;16(3):282–293.

doi:10.1080/15402002.2016.1210147

34. Mindell JA, Cook RA,

Nikolovski J. Sleep patterns and sleep disturbances across pregnancy. *Sleep Med.* 2015;16(4):483–488. doi:10.1016/j.sleep.2014.12.006

35.

Mitchell-Jones N, Gallos I, Farren J, Tobias A, Bottomley C, Bourne T. Psychological morbidity associated with hyperemesis gravidarum: a systematic review and meta-analysis. *BJOG.* 2017;124(1):20–30.

doi:10.1111/1471-0528.14180

36. Hoirisch-Clapauch S, Brenner B, Nardi AE. Adverse obstetric and neonatal outcomes in women with mental disorders. *Thromb Res.* 2015;135:S60–S63. doi:10.1016/S0049-3848(15)50446-5

37. Chou FH, Avant KC, Kuo SH, Fetzer SJ.

Relationships between nausea and vomiting, perceived stress, social support, pregnancy planning, and psychosocial adaptation in a sample of mothers: a questionnaire survey. *Int J Nurs Stud.* 2008;45(8):1185–1191. doi:10.1016/j.ijnurstu.2007.08.00

38. Lönnberg G, Jonas W, Unternaehrer E, Bränström R, Nissen E, Niemi M. Effects of a mindfulness based childbirth and parenting program on pregnant

women's perceived stress and risk of perinatal depression-Results from a randomized controlled trial. *J Affect Disord.* 2020;262:133–142. doi:10.1016/j.jad.2019.10.048

39. Rodríguez MA, Valentine J, Ahmed SR, et al. Intimate partner violence and maternal depression during the perinatal period: a longitudinal investigation of Latinas. *Violence Against Women.* 2010;16(5):543–559. doi:10.1177/1077801210366959

40. Keller J, Gomez R, Williams G, et al. HPA axis in major depression: cortisol,