

First record of *Plagiorchis muris* Tanabe, 1922 from black rat *Rattus rattus* in Qarmat Ali, Basrah, Iraq

Khalid. J. K. Al-Zihiry* A. H. H. Awad**

*Department of Microbiology/ College of Medicine/ University of Thi-qar

**Department of Biology / College of Education University of Basrah

Abstract:

During the period from March till May 2004 a total of 20 black rats *Rattus rattus* were collected from Qarmat Ali ,Basrah .About 15% of these rats were found infected with digenetic trematode *Plagiorchis muris*. Full description ,measurements and a comparison with other studies in the world were given.The presence of this trematode in the present study is considered a first record in Iraq.

Introduction:

The rodents, particularly those living in close association with man , play a significant role of human health (Stojcevic,*et al.*,2004); its act as definitive and/or intermediate hosts of many endoparasites including helminthes with zoonotic potential (Ito and Itagaki,2003).Many of studies were carried out on rodent parasites in the world ,in contrast a few of those were carried out in different parts of Iraq. Two studies were done in Basrah province, Al-Hadithi *et al.*(1985) collect 50 of Norway rats *Rattus norvegicus* from center of Basrah and five helminthes recorded, three of cestode and two of nematode. Al-Zihiry (2002) collect 450 small mammals from many localities in Basrah province and he record two of trematodes from genus *Echinostoma* in black rats *R.rattus* and *R.norvegicus*. The aim of present study was to identify the main helminthes of *R.rattus* in Qarmat Ali, Basrah.

Materials and Methods:

A total of 20 specimens of black rats *Rattus rattus* (Rodentia) were collected from Qarmat Ali, Basrah during the period from March till May 2004 using number of live traps. The animals were dissected and their guts were opened in

petridishes contain normal saline and examined for helminthes on dissecting microscope. The recovered trematodes were flattened, fixed in 70% ethanol, stained with semichon acid carmine, dehydrated in ethanol, cleared in toullene and mounted on clean slides with Canada balsam (Garcia and Ash,1979).The measurements were taken by ocular micrometer of Olympus microscope and adult trematodes were drawn by camera lucida.Yamaguti (1958) and Seo *et al.*(1964) were depended to identify these trematodes.

Results:

Eighteen specimens of *Plagiorchis muris* Tanabe, 1922 were isolated from the small intestine of three 15% black rat *Rattus rattus* with intensity of infection equal to 6.

Description : (All measurement was in millimeters which based on three specimens)

Fig.1

Body elongate with spinose cuticle in interior half of body, 1.029 to 1.784 long by 0.315 to 0.415 wide in widest area at the level of ventral sucker. Oral sucker spherical, 0.124 to 0.166 in diameter. Pharynx muscular, 0.060 to 0.075 long by 0.042 to 0.060 wide. Esophagus short, bifurcation in two branches(Ceca)extending to posterior end of the body. Ventral sucker spherical, 0.100 to 0.130 in diameter and located in one third of body at the distance 0.350 to 0.550 from interior end of body. Testes oval, tandem, lying in posterior half of body. Interior testis at the right side of median line, 0.182 to 0.200 long by 0.150 to 0.160 wide. Posterior testis at the left side of median line, 0.180 to 0.210 long by 0.140 to 0.152. Cirrus sac elongate, located between ventral sucker and cecal branches and extends to level of ovary, 0.300 to 0.420 long by 0.060 to 0.080 wide.

Seminal vesicle is lying in posterior part of cirrus sac. Ovary spherical, located in the right side of median line beyond the ventral sucker, 0.112 to 0.130 in diameter. Vitteline glands follicular, lateral, extending from the pharynx level to posterior end of body. Uterus is in (S) shape and located in restricted area between end of ventral sucker and level of posterior testis. Egg oval, 0.030 to 0.032 long by 0.014 to 0.020 wide.

Host: *Rattus rattus*

Location: small intestine

Locality: Qarmat Ali, Basrah




Fig. (1) : Adult of *Plagiorchis muris* , ventral view

Discussion:

The finding of trematode *plagiorchis muris* in the present study is considered as a first record in Iraq, while its recorded by Seo *et al.*,(1964) from *Apodemus agrarius* and *Rattus rattus* in Korea, its isolated in Taiwan by Fischthal and Kuntz (1975) from small intestine of *R. rattus*. In India Gupta and Jahan (1978) isolate the trematode from small intestine of *Crocidura* sp., through the study carried out in urban, rural residential and other ecological areas of Nigeria, these trematode was collected from *R.rattus* (Udonsi,1989), moreover its reported by Ito and Itagaki (2003) parasitizing in small intestine of *A.speciosus* in Japan. The measurments of the present specimens are similar to those taken by Seo *et al.*,(1964) and Ishimoto (1974) with some differences but its different from measurments of the specimens isolated by Gupta and Jahan (1978) from *Crocidura* sp. (tab.1). These differences

can be attributed to the abnormal host (*Crocidura* sp.) to these trematode. Tanabe (1922) was the first who described this trematode in both *R. rattus* and *R. norvegicus* of Kyoto, Japan and he reports that the snails *Lymnaea pervica* and *Stagnicola emarginata* were served as a first intermediate hosts. The second intermediate hosts are aquatic insects, such as mosquito larvae, insect naiads, freshwater snails and freshwater fishes (Asada *et al.*, 1962; Komiya, 1965). Recently, six species of dragonflies were reported that infected with *P. muris* metacercariae and detected as a second intermediate hosts for this trematode (Hong *et al.*, 1999). *P. muris* is the unique member of the family Plagiorchiidae is known as zoonotic pathogen by human infection (Lee *et al.*, 2004). Several cases of human infection by this trematode were reported in parts of world, especially in Asia, from these infections, 500 cases were recorded in Korea (Hong *et al.*, 1996; WHO, 2002). Experimental infections were carried out with these trematode on human in Michigan when 150 metacercariae were ingested. On the ninth day after the initial infection the first eggs (0.038 x 0.019) in morula stage were found in the stool, the egg production was reached to 74,700 egg per day on the seventeenth day after infection, the pathological nature were detected in these studies (McMullen, 1936).

Table (1): The comparison measurements of *plagiorchis muris* in present study and other studies

	Present study <i>Rattus rattus</i>	Seo <i>et al.</i> , 1964 <i>R. rattus</i> , <i>A. agrarius</i>	Ishimoto, 1974 <i>Clethrionomys</i> <i>Apodemus</i> <i>A. speciosus</i>	<i>rufocanus</i> , <i>argenteus</i> ,	Gupta and Jahan, 1978 <i>Crocidura</i> sp.
Body					
Long	1.029-1.784	1.460-1.770	1.40-1.89		2.870
Wide	0.315-0.415	0.364-0.448	0.38-0.51		0.900
Oral sucker					
Long	0.124-0.166	0.154-0.168	0.147-0.207		0.220
Wide	in diameter	in diameter	in diameter		0.270
Pharynx					
Long	0.060-0.075	0.066-0.084	0.079-0.105		0.160
Wide	0.042-0.060	0.076-0.084	0.058-0.084		in diameter
Ventral sucker					
	0.100-0.130 in diameter	0.154-0.168 in diameter	0.126-0.172 in diameter		0.330 in diameter
Location of ventral sucker from interior end	0.350-0.550	0.420-0.644	One third		0.700
Cirrus sac					
Long	0.300-0.420	0.350-0.396	0.51-0.60		0.550
Wide	0.060-0.080	0.059-0.070	0.089-0.092		0.060
Interior testis					
Long	0.182-0.200	0.182-0.266	0.14-0.24		-----
Wide	0.150-0.160	0.168-0.238	in diameter		
Posterior testis					
Long	0.180-0.210	0.224-0.280	0.14-0.24		0.520

Wide	0.140-0.152	0.154-0.210	in diameter	0.460
Ovary				
Long	0.112-0.130	0.154-0.210	0.10-0.18	0.240
Wide	in diameter	0.154-0.196	in diameter	0.280
Eggs				
Long	0.030-0.032	0.030-0.033	0.033	0.030-0.040
Wide	0.014-0.020	0.020-0.023	0.021	0.012-0.020

References:

- Al-Hadithi, I.A.W.; Habash, A.H. and Alwan, M.J.(1985). The incidence of helminth parasites of the digestive tract of *Rattus norvegicus* from Basrah area, Iraq . J. Biol. Sci. Res.,16:137-154.
- Al-Zihiry, K.J.K.(2002).Helminth parasites in some small mammals from Basrah province .M.Sc.thesis, Coll.Edu., Univ.Basrah:84 pp.
- Asada, J.I.; Otagaki, H.; Morita, M.; Takeuchi, T.; Sakai, Y.; Konishi, T. and Okahashi, K.(1962).A case report on the human infection with *Plagiorchis muris* Tanabe, 1922 in Japan.Jpn.J parasitol., 1: 512-516.
- Fischthal, J.H and Kuntz, R.E.(1975).Some digenetic trematode of mammals from Taiwan .Proc.Helminthol.Soc.Washington,42:149-157.
- Garcia L.S.and Ash,L.R.(1979).Diagnostic parasitology clinical laboratory manual.2nd edn.,The C.V.Mosby company st.Louis,174pp.
- Gupta,V. and Jahan,A.(1978). On a new trematode, *Plagiorchis funambulusi* sp.nov. and *Plagiorchis muris* Tanabe, 1922 Digenea:Plagiorchidae(Luhe, 1901) Ward, 1917 from mammals.Indian J.Helminthol.,30(2):143-150.
- Hong, S.J.; Woo, H.C.; Chai, J.Y.(1996). A human case of *Plagiorchis muris* (Tanabe, 1922: Digenea) in the Republic of Korea: Freshwater fishes as a possible source of infection. J. Parasitol. 82: 647-649.
- Hong, S.J.; Woo, H.C.; Lee, S.U.; Huh, S.(1999). Infection status of dragonflies with *Plagiorchis muris* metacercariae in Korea. Korean J.Parasitol.,37(2):65-70.
- Ishimoto, Y.(1974). Studies on helminthes of voles in Hokkaido I. Taxonomical study. Jap. J. Vet. Res., 22: 1-12.
- Ito, M. and Itagaki,T.(2003).Survey on wild rodents for endoparasites in Iwate prefecture, Japan.J.Vet.Med.Sci.,65(10):1151-1153.
- Komiya, Y.(1965). Metacercariae in Japan and adjacent territories.In progress of Medical Parasitology in Japan.Vol. II. pp.225-233, Meguro Parasitological Museum, Tokyo, Japan.
- Lee,S.; Huh,S. and Sohn,W.(2004). Molecular phylogenic location of the *Plagiorchis muris* (Digenea: Plagiorchidae) based on sequences of partial 28 S D1 rDNA and mitochondrial cytochrome C oxidase subunit I.Korean J.Parasitol., 42(1):71-75.
- McMullen,D.B.(1936).An experimental infection of *Plagiorchis muris* in man.J.Parasitol.,22:113-115.
- Seo, B.S.;Rim, H.J. and Lee,C.W.(1964).Studies on the parasitic helminthes of Korea I. Trematodes of rodents.Korean J.Parasitol., 2(1):20-26.

Stojcevic, D.;Mihaljevic,Z.and Marinculic,A.(2004).Parasitological survey of rats in rural regions of Croatia.Vet.Med.Czech,3:70-74.

Tanabe,H.(1922).Contributions to the knowledge of the developmental cycle of digenetic trematodes.On a new species of trematode *Lepoderma muris* n.sp.Okayama Igakkai Zasshi,385:47-57 (in Japanese).

Udonsi,J.K.(1989).Helminth parasites of wild populations of the black rat, *Rattus rattus* (L.), from urban, rural residential and other ecological areas of Nigeria.Acta parasitol.Pol., 34:107-116.

WHO (2002).Food-borne trematode infection in Asia.WHO/FAO Workshop.Rep. Ser. Geneva, No.40.

Yamaguti, S.(1958).Systema helminthum,I: The trematode of vertebrates. Part I+II.Intersci. publ.Inc.Ltd.New York.

تسجيل جديد للمثقوبة *plagiorchis muris* من الجرذ الأسود *Rattus rattus*

في منطقة كرمة علي ، البصرة ، العراق

خالد جميل كاظم الزهيري * عبد الحسين حبش عواد**

*فرع الأحياء المجهرية/ كلية الطب/ جامعة ذي قار

**قسم علوم الحياة / كلية التربية / جامعة البصرة

الخلاصة

جمع 20 جرذ اسود *Rattus rattus* خلال الفترة من شهر اذار ولغاية نهاية شهر ايار عام 2004 من منطقة كرمة علي في محافظة البصرة وقد وجد ان 15% منها مصابة بالمثقوبة ثنائية المنشأ *Plagiorchis muris*. أعطى الوصف والقياسات الكاملة للمثقوبة وعملت مقارنة لقياسات الدودة المعزولة مع اخرى في دراسات سابقة اجريت في العالم . يعد وجود هذه المثقوبة في الدراسة الحالية هو التسجيل الأول لها في العراق .